

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 March 2002 (21.03.2002)

PCT

(10) International Publication Number
WO 02/22636 A1

(51) International Patent Classification⁷: C07H 21/02, 21/04, A61K 48/00, C12N 15/85, 15/86, C12Q 1/68, C12P 19/34

(21) International Application Number: PCT/US01/28572

(22) International Filing Date:
12 September 2001 (12.09.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/663,834 15 September 2000 (15.09.2000) US

(71) Applicant (for all designated States except US): **ISIS PHARMACEUTICALS, INC.** [US/US]; 2292 Faraday Avenue, Carlsbad, CA 92008 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BENNETT, C., Frank** [US/US]; 1347 Cassins Street, Carlsbad, CA 92008 (US). **COWSERT, Lex, M.** [US/US]; 1299 Parrott Drive, San Mateo, CA 94402 (US).

(74) Agents: **LICATA, Jane, Massey et al.**; Licata & Tyrell P.C., 66 E. Main Street, Marlton, NJ 08053 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/22636 A1

(54) Title: ANTISENSE MODULATION OF HER-2 EXPRESSION

(57) Abstract: Antisense compounds, compositions and methods are provided for modulating the expression of Her-2. The compositions comprise antisense compounds, particularly antisense oligonucleotides, targeted to nucleic acids encoding Her-2. Methods of using these compounds for modulation of Her-2 expression and for treatment of diseases associated with expression of Her-2 are provided.

-1-

ANTISENSE MODULATION OF HER-2 EXPRESSION

5

FIELD OF THE INVENTION

The present invention provides compositions and
10 methods for modulating the expression of Her-2. In
particular, this invention relates to compounds,
particularly oligonucleotides, specifically hybridizable
with nucleic acids encoding Her-2. Such oligonucleotides
have been shown to modulate the expression of Her-2.

15

BACKGROUND OF THE INVENTION

One of the principal mechanisms by which cellular
regulation is effected is through the transduction of
extracellular signals into intracellular signals that in
20 turn modulate biochemical pathways. Examples of such
extracellular signaling molecules include growth factors,
cytokines, and chemokines. The cell surface receptors of
these molecules and their associated signal transduction
pathways are therefore one of the principal means by which
25 cellular behavior is regulated. Because cellular
phenotypes are largely influenced by the activity of these
pathways, it is currently believed that a number of disease
states and/or disorders are a result of either aberrant
activation or functional mutations in the molecular
30 components of signal transduction pathways. Consequently,
considerable attention has been devoted to the
characterization of these receptor proteins.

HER2 (also known as Human EGF Receptor 2, c-erbB2,
erbB2 and Neu), a member of the EGF family of

-2-

receptor/tyrosine kinases, is a protein that has been shown to play a complex role in several signal transduction pathways by forming homo- and heterodimers with other members of the EGF family depending on their concentrations and the concentration of particular ligands. The ligands for the EGF family of receptors include six ligands that bind EGFR (ErbB1) including EGF, TGF-alpha, amphiregulin, heparin binding EGF-like growth factor, betacellulin and epiregulin. In addition, there are two groups of ligands specific to HER3 and HER4 collectively termed neuregulins (NRGs) because of their demonstrated role in the nervous system. Although HER2 acts as the most oncogenic member of the family by inhibition ligand dissociation from the dimer, there is no known ligand specific to HER2. HER3 and HER4 function as the binding receptors; and HER2 along with EGFR are considered co-receptors and are recruited as partners to HER3 and HER4 upon ligand binding. (Burden and Yarden, *Neuron*, 1997, 18, 847-855).

HER2 was first isolated and cloned in 1985 from brain tumors of female mice treated with a carcinogen which resulted in an oncogenic mutation of the receptor. This mutation transformed the receptor from a ligand dependent kinase to a constitutively active kinase which, when expressed early in development, caused brain tumors (Earp et al., *Breast Cancer Res. Treat.*, 1995, 35, 115-132). Since then it has been demonstrated that HER2 plays a role in both neural and cardiac development (Carraway, *BioEssays*, 1996, 18, 263-266).

Cellular transformation and acquisition of the metastatic phenotype are the two main changes normal cells undergo during the progression to cancer. HER2 has been implicated in both processes. The HER2 receptor has been shown to be altered in a number of malignancies including

-3-

gastric, esophageal, salivary, colon, bladder and lung cancer and overexpression of HER2 in advanced breast cancer has been correlated with the progression of the disease (Earp et al., *Breast Cancer Res. Treat.*, 1995, 35, 115-
5 132).

Currently, there are no known therapeutic agents which effectively inhibit the synthesis of HER2. Consequently, there remains a long felt need for additional agents capable of effectively inhibiting HER2 expression and/or
10 function. To date, strategies aimed at inhibiting HER2 function have involved the use of gene knockouts in mice, receptor antibodies, antisense vectors and oligonucleotides, ribozymes and antigen-activated cytotoxic lymphocytes (Cirisano and Karlan, *J. Soc. Gynecol.*
15 *Investig.*, 1996, 3, 99-105).

A critical role for HER2 during gestational development was illustrated using HER2 deficient mice which underwent aberrant cardiac development 10 days post-fertilization and died (Alroy and Yarden, *FEBS Lett.*, 1997,
20 410, 83-86). Other studies demonstrated the importance of HER2 in neural development. Mice lacking in either NRG or HER2 expression lacked neural crest-derived cells of cranial sensory ganglia as well as axonal connections to the hindbrain (Carraway, *BioEssays*, 1996, 18, 263-266).

Other studies using an expression vector encoding HER2 in the antisense orientation have shown that the reduction of HER2 levels in various cancer cells may have a beneficial effect. For example, Yoo and Hamburger demonstrated that HER2 expression modulates signal
25 transduction pathways in breast cancer lines (Yoo and Hamburger, *Mol. Cell. Endocrinol.*, 1998, 138, 163-171), while Pegues and Stromberg demonstrated that inhibition of HER2 reduced anchorage independent growth of ovarian

-4-

carcinoma cells (Pegues and Stromberg, *Cancer Letters*, 1997, 117, 73-79). Finally, Casalini et al. have shown that HER2 antisense constructs inhibit tumorigenicity in lung adenocarcinoma cells (Casalini et al., *Int. J. Cancer*, 5 1997, 72, 631-636).

Furthermore, Herceptin™, a recombinant humanized anti-HER2 antibody has been approved for treatment of metastatic breast cancer and has been shown to enhance antitumor activity of Paclitaxel and Doxorubicin in breast cancer xenografts (Baselga et al., *Cancer Res.*, 1998, 58, 2825-10 2831).

Other studies of antisense oligonucleotides targeting HER2 in breast carcinoma cells also demonstrated growth inhibition of the cells using an oligonucleotide targeting the 5' region of the gene and synergistic effects were seen 15 with treatment of the oligonucleotide in combination with conventional chemotherapeutic agents (Roh et al., *Surgery*, 1999, 126, 413-421; Roh et al., *J. Surg. Res.*, 1998, 77, 85-90). The same oligonucleotide was subsequently shown to induce apoptosis in cancer cells overexpressing HER2 (Roh 20 et al., *Cancer Res.*, 2000, 60, 560-565). Several studies have also shown the utility of antisense technology in downregulating HER2 expression in cancer cells (Bennett et al., 1999; Bertram et al., *Biochemical and Biophysical Research Communications*, 1994, 2000, 661-667; Liu and Pogo, 25 *Antisense and Nucleic Acid Drug Development*, 1996, 6, 9-16; Marks et al., 1998; Vaughn et al., *Proc. Natl. Acad. Sci. U. S. A.*, 1995, 92, 8338-8342). Finally, Urbaine et al. showed that cells expressing a certain mRNA preferentially 30 retain the antisense probe targeting that mRNA. These studies concluded that antisense molecules represent a novel class of radiopharmaceutical drugs with unique

-5-

specificity (Urbain et al., *Eur. J. Nucl. Med.*, 1995, 22, 499-504).

Disclosed in the PCT Publication WO 99/31118 are recombinant adenoviruses which express ribozymes targeting
5 HER2 mRNA for the purpose of inhibiting tumor growth (Czubayko and Wellstein, 1999). Ribozymes are also disclosed in Wiechen et al. for use in downregulating HER2 in ovarian cancers (Wiechen et al., *Cancer Gene Ther.*, 1998, 5, 45-51).

10 Antisense technology is emerging as an effective means for reducing the expression of specific gene products and may therefore prove to be uniquely useful in a number of therapeutic, diagnostic, and research applications for the modulation of HER2 expression.

15 The present invention provides compositions and methods for modulating HER2 expression.

SUMMARY OF THE INVENTION

The present invention is directed to compounds,
20 particularly antisense compounds, which are targeted to a nucleic acid encoding Her-2, and which modulate the expression of Her-2. Pharmaceutical and other compositions comprising the antisense compounds of the invention are also provided. Further provided are methods of modulating
25 the expression of Her-2 in cells or tissues comprising contacting said cells or tissues with one or more of the antisense compounds or compositions of the invention. Further provided are methods of treating an animal, particularly a human, suspected of having or being prone to
30 a disease or condition associated with expression of Her-2 by administering a therapeutically or prophylactically effective amount of one or more of the antisense compounds or compositions of the invention.

-6-

DETAILED DESCRIPTION OF THE INVENTION

The present invention employs oligomeric antisense compounds, particularly oligonucleotides, for use in modulating the function of nucleic acid molecules encoding Her-2, ultimately modulating the amount of Her-2 produced.

This is accomplished by providing antisense compounds which specifically hybridize with one or more nucleic acids encoding Her-2. As used herein, the terms "target nucleic acid" and "nucleic acid encoding Her-2" encompass DNA encoding Her-2, RNA (including pre-mRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as "antisense". The functions of DNA to be interfered with include replication and transcription. The functions of RNA to be interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA.

The overall effect of such interference with target nucleic acid function is modulation of the expression of Her-2. In the context of the present invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene. In the context of the present invention, inhibition is the preferred form of modulation of gene expression and mRNA is a preferred target.

-7-

It is preferred to target specific nucleic acids for antisense. "Targeting" an antisense compound to a particular nucleic acid, in the context of this invention, is a multistep process. The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target is a nucleic acid molecule encoding Her-2. The targeting process also includes determination of a site or sites within this gene for the antisense interaction to occur such that the desired effect, e.g., detection or modulation of expression of the protein, will result. Within the context of the present invention, a preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function *in vivo*. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be

-8-

preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the
5 codon or codons that are used *in vivo* to initiate translation of an mRNA molecule transcribed from a gene encoding Her-2, regardless of the sequence(s) of such codons.

It is also known in the art that a translation
10 termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of
15 such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene
20 that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon.

The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the
25 translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the
30 translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), known in

-9-

the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap.

10 The 5' cap region may also be a preferred target region.

Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. mRNA splice sites, i.e., intron-exon junctions, may also be preferred target regions, and are particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also preferred targets. It has also been found that introns can also be effective, and therefore preferred, target regions for antisense compounds targeted, for example, to DNA or pre-mRNA.

15
20
25

Once one or more target sites have been identified, oligonucleotides are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

30

In the context of this invention, "hybridization" means hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between

-10-

complementary nucleoside or nucleotide bases. For example, adenine and thymine are complementary nucleobases which pair through the formation of hydrogen bonds.

"Complementary," as used herein, refers to the capacity for
5 precise pairing between two nucleotides. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to
10 each other at that position. The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable"
15 and "complementary" are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is understood in the art that the sequence of an antisense
20 compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. An antisense compound is specifically hybridizable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or
25 RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of *in vivo*
30 assays or therapeutic treatment, and in the case of *in vitro* assays, under conditions in which the assays are performed.

-11-

Antisense and other compounds of the invention which hybridize to the target and inhibit expression of the target are identified through experimentation, and the sequences of these compounds are hereinbelow identified as preferred embodiments of the invention. The target sites to which these preferred sequences are complementary are hereinbelow referred to as "active sites" and are therefore preferred sites for targeting. Therefore another embodiment of the invention encompasses compounds which hybridize to these active sites.

Antisense compounds are commonly used as research reagents and diagnostics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes. Antisense compounds are also used, for example, to distinguish between functions of various members of a biological pathway. Antisense modulation has, therefore, been harnessed for research use.

The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses. Antisense oligonucleotide drugs, including ribozymes, have been employed as therapeutic moieties in the treatment of disease states in animals and man. Antisense oligonucleotides have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic modalities that can be configured to be useful in treatment regimes for treatment of cells, tissues and animals, especially humans. In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of

-12-

ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as
5 oligonucleotides having non-naturally-occurring portions which function similarly. Such modified or substituted oligonucleotides are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic
10 acid target and increased stability in the presence of nucleases.

While antisense oligonucleotides are a preferred form of antisense compound, the present invention comprehends other oligomeric antisense compounds, including but not
15 limited to oligonucleotide mimetics such as are described below. The antisense compounds in accordance with this invention preferably comprise from about 8 to about 50 nucleobases (i.e. from about 8 to about 50 linked nucleosides). Particularly preferred antisense compounds
20 are antisense oligonucleotides, even more preferably those comprising from about 12 to about 30 nucleobases. Antisense compounds include ribozymes, external guide sequences (EGS) oligonucleotides (oligozymes), and other short catalytic RNAs or catalytic oligonucleotides which
25 hybridize to the target nucleic acid and modulate its expression.

As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes
30 of such heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that

-13-

include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

Specific examples of preferred antisense compounds useful in this invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and

-14-

those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included.

5 Representative United States patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S.: 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 10 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

15 Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short 20 chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; 25 methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH₂ component parts.

30 Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S.: 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564;

-15-

5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677;
5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289;
5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070;
5,663,312; 5,633,360; 5,677,437; and 5,677,439, certain of
5 which are commonly owned with this application, and each of
which is herein incorporated by reference.

In other preferred oligonucleotide mimetics, both the
sugar and the internucleoside linkage, i.e., the backbone,
of the nucleotide units are replaced with novel groups.
10 The base units are maintained for hybridization with an
appropriate nucleic acid target compound. One such
oligomeric compound, an oligonucleotide mimetic that has
been shown to have excellent hybridization properties, is
referred to as a peptide nucleic acid (PNA). In PNA
15 compounds, the sugar-backbone of an oligonucleotide is
replaced with an amide containing backbone, in particular
an aminoethylglycine backbone. The nucleobases are
retained and are bound directly or indirectly to aza
nitrogen atoms of the amide portion of the backbone.
20 Representative United States patents that teach the
preparation of PNA compounds include, but are not limited
to, U.S.: 5,539,082; 5,714,331; and 5,719,262, each of
which is herein incorporated by reference. Further teaching
of PNA compounds can be found in Nielsen et al., *Science*,
25 1991, 254, 1497-1500.

Most preferred embodiments of the invention are
oligonucleotides with phosphorothioate backbones and
oligonucleosides with heteroatom backbones, and in
particular -CH₂-NH-O-CH₂-, -CH₂-N(CH₃)-O-CH₂- [known as a
30 methylene (methylimino) or MMI backbone], -CH₂-O-N(CH₃)
-CH₂-, -CH₂-N(CH₃)-N(CH₃)-CH₂- and -O-N(CH₃)-CH₂-CH₂- [wherein
the native phosphodiester backbone is represented as -O-P-
O-CH₂-] of the above referenced U.S. patent 5,489,677, and

-16-

the amide backbones of the above referenced U.S. patent 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced U.S. patent 5,034,506.

5 Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and
10 alkynyl may be substituted or unsubstituted C₁ to C₁₀ alkyl or C₂ to C₁₀ alkenyl and alkynyl. Particularly preferred are O[(CH₂)_nO]_mCH₃, O(CH₂)_nOCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, and O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10. Other preferred oligonucleotides
15 comprise one of the following at the 2' position: C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino,
20 substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred
25 modification includes 2'-methoxyethoxy (2'-O-CH₂CH₂OCH₃, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta*, 1995, 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification includes 2'-dimethylaminoethoxy, i.e., a O(CH₂)₂ON(CH₃)₂
30 group, also known as 2'-DMAOE, as described in examples hereinbelow, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE),

-17-

i.e., 2'-O-CH₂-O-CH₂-N(CH₂)₂, also described in examples hereinbelow.

Other preferred modifications include 2'-methoxy (2'-O-CH₃), 2'-aminopropoxy (2'-OCH₂CH₂CH₂NH₂) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S.: 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil

-18-

(pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further nucleobases include those disclosed in United States Patent No. 3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., *Angewandte Chemie*, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y.S., Chapter 15, *Antisense Research and Applications*, pages 289-302, Crooke, S.T. and Lebleu, B., ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

Representative United States patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. 3,687,808, as well as U.S.: 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908;

-19-

5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121,
5,596,091; 5,614,617; and 5,681,941, certain of which are
commonly owned with the instant application, and each of
which is herein incorporated by reference, and United
5 States patent 5,750,692, which is commonly owned with the
instant application and also herein incorporated by
reference.

Another modification of the oligonucleotides of the
invention involves chemically linking to the
10 oligonucleotide one or more moieties or conjugates which
enhance the activity, cellular distribution or cellular
uptake of the oligonucleotide. Such moieties include but
are not limited to lipid moieties such as a cholesterol
moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 1989,
15 86, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med.*
Chem. Lett., 1994, 4, 1053-1060), a thioether, e.g., hexyl-
S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*,
1992, 660, 306-309; Manoharan et al., *Bioorg. Med. Chem.*
Let., 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et
20 al., *Nucl. Acids Res.*, 1992, 20, 533-538), an aliphatic
chain, e.g., dodecandiol or undecyl residues (Saison-
Behmoaras et al., *EMBO J.*, 1991, 10, 1111-1118; Kabanov et
al., *FEBS Lett.*, 1990, 259, 327-330; Svinarchuk et al.,
Biochimie, 1993, 75, 49-54), a phospholipid, e.g., di-
25 hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-
hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al.,
Tetrahedron Lett., 1995, 36, 3651-3654; Shea et al., *Nucl.*
Acids Res., 1990, 18, 3777-3783), a polyamine or a
polyethylene glycol chain (Manoharan et al., *Nucleosides &*
30 *Nucleotides*, 1995, 14, 969-973), or adamantane acetic acid
(Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651-3654),
a palmityl moiety (Mishra et al., *Biochim. Biophys. Acta*,
1995, 1264, 229-237), or an octadecylamine or hexylamino-

-20-

carbonyl-oxycholesterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.*, 1996, 277, 923-937.

Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but
5 are not limited to, U.S.: 4,828,979; 4,948,882; 5,218,105;
5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717,
5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802;
5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718;
5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779;
10 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582;
4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830;
5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506;
5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241,
5,391,723; 5,416,203, 5,451,463; 5,510,475; 5,512,667;
15 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481;
5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and
5,688,941, certain of which are commonly owned with the
instant application, and each of which is herein
incorporated by reference.

20 It is not necessary for all positions in a given
compound to be uniformly modified, and in fact more than
one of the aforementioned modifications may be incorporated
in a single compound or even at a single nucleoside within
an oligonucleotide. The present invention also includes
25 antisense compounds which are chimeric compounds.
"Chimeric" antisense compounds or "chimeras," in the
context of this invention, are antisense compounds,
particularly oligonucleotides, which contain two or more
chemically distinct regions, each made up of at least one
30 monomer unit, i.e., a nucleotide in the case of an
oligonucleotide compound. These oligonucleotides typically
contain at least one region wherein the oligonucleotide is
modified so as to confer upon the oligonucleotide increased

-21-

resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. An additional region of the oligonucleotide may serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of oligonucleotide inhibition of gene expression. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

Chimeric antisense compounds of the invention may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described above. Such compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S.: 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

The antisense compounds used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis.

-22-

Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, CA). Any other means for such synthesis known in the art may additionally or alternatively be employed. It is well
5 known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and alkylated derivatives.

The antisense compounds of the invention are synthesized *in vitro* and do not include antisense compositions of biological origin, or genetic vector
10 constructs designed to direct the *in vivo* synthesis of antisense molecules.

The compounds of the invention may also be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as
15 for example, liposomes, receptor targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption assisting
20 formulations include, but are not limited to, U.S.:
5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291;
5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330;
4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170;
5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978;
25 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259;
5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of which is herein incorporated by reference.

The antisense compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such
30 esters, or any other compound which, upon administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the

-23-

disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the invention, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents.

5 The term "prodrug" indicates a therapeutic agent that is prepared in an inactive form that is converted to an active form (i.e., drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or conditions. In particular, prodrug versions of the
10 oligonucleotides of the invention are prepared as SATE [(S-acetyl-2-thioethyl) phosphate] derivatives according to the methods disclosed in WO 93/24510 to Gosselin et al., published December 9, 1993 or in WO 94/26764 and U.S. 5,770,713 to Imbach et al.

15 The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto.

20 Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are
25 N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge et al., "Pharmaceutical Salts," *J. of Pharma Sci.*, 1977, 66, 1-19). The base addition salts of said acidic compounds
30 are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and

-24-

isolating the free acid in the conventional manner. The free acid forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention. As used herein, a "pharmaceutical addition salt" includes a pharmaceutically acceptable salt of an acid form of one of the components of the compositions of the invention. These include organic or inorganic acid salts of the amines. Preferred acid salts are the hydrochlorides, acetates, salicylates, nitrates and phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid, glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic acid; and with amino acids, such as the 20 alpha-amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic

-25-

acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid. Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible.

For oligonucleotides, preferred examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine.

The antisense compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. For therapeutics, an animal, preferably a human, suspected of having a disease or disorder which can be treated by modulating the expression of Her-2 is treated by administering antisense

-26-

compounds in accordance with this invention. The compounds of the invention can be utilized in pharmaceutical compositions by adding an effective amount of an antisense compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the antisense compounds and methods of the invention may also be useful prophylactically, e.g., to prevent or delay infection, inflammation or tumor formation, for example.

The antisense compounds of the invention are useful for research and diagnostics, because these compounds hybridize to nucleic acids encoding Her-2, enabling sandwich and other assays to easily be constructed to exploit this fact. Hybridization of the antisense oligonucleotides of the invention with a nucleic acid encoding Her-2 can be detected by means known in the art. Such means may include conjugation of an enzyme to the oligonucleotide, radiolabelling of the oligonucleotide or any other suitable detection means. Kits using such detection means for detecting the level of Her-2 in a sample may also be prepared.

The present invention also includes pharmaceutical compositions and formulations which include the antisense compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial,

-27-

subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration.

Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful.

Compositions and formulations for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, sachets or tablets. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable.

Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions may be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids.

The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such

-28-

techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

In one embodiment of the present invention the pharmaceutical compositions may be formulated and used as foams. Pharmaceutical foams include formulations such as, but not limited to, emulsions, microemulsions, creams, jellies and liposomes. While basically similar in nature these formulations vary in the components and the consistency of the final product. The preparation of such compositions and formulations is generally known to those skilled in the pharmaceutical and formulation arts and may be applied to the formulation of the compositions of the present invention.

30 Emulsions

The compositions of the present invention may be prepared and formulated as emulsions. Emulsions are typically heterogenous systems of one liquid dispersed in

-29-

another in the form of droplets usually exceeding 0.1 μm in diameter. (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in

5 *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in

10 *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 301). Emulsions are often biphasic systems comprising of two immiscible liquid phases intimately mixed and dispersed with each other. In

15 general, emulsions may be either water-in-oil (w/o) or of the oil-in-water (o/w) variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase the resulting composition is called a

20 water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase the resulting composition is called an oil-in-water (o/w) emulsion.

25 Emulsions may contain additional components in addition to the dispersed phases and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical

30 excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed. Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and

water-in-oil-in-water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which

-30-

individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water stabilized in an oily continuous provides an o/w/o
5 emulsion.

Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into the external or continuous phase and maintained in this
10 form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of stabilizing emulsions entail the use of emulsifiers that may be
15 incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker
20 (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (Rieger,
25 in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., 1988, volume 1, p. 199). Surfactants are
30 typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool

-31-

in categorizing and selecting surfactants in the preparation of formulations. Surfactants may be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and
5 amphoteric (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides,
10 lecithin and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers
15 especially in combination with surfactants and in viscous preparations. These include polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as bentonite, attapulgite, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium
20 aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes,
25 fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants (Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in *Pharmaceutical Dosage
30 Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Hydrophilic colloids or hydrocolloids include naturally occurring gums and synthetic polymers such as

-32-

polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth), cellulose derivatives (for example, carboxymethylcellulose and carboxypropylcellulose), and synthetic polymers (for
5 example, carbomers, cellulose ethers, and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming strong interfacial films around the dispersed-phase droplets and by increasing the viscosity of the external
10 phase.

Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides that may readily support the growth of microbes, these formulations often incorporate preservatives. Commonly
15 used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added to emulsion formulations to prevent deterioration of the
20 formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid,
25 and lecithin.

The application of emulsion formulations via dermatological, oral and parenteral routes and methods for their manufacture have been reviewed in the literature (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger
30 and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral delivery have been very widely used because of reasons of ease of formulation, efficacy from an absorption and

-33-

bioavailability standpoint. (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

10 In one embodiment of the present invention, the compositions of oligonucleotides and nucleic acids are formulated as microemulsions. A microemulsion may be defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable
15 liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and
20 then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are
25 stabilized by interfacial films of surface-active molecules (Leung and Shah, in: *Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215). Microemulsions commonly are prepared via a combination of three to five
30 components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used

-34-

and on the structure and geometric packing of the polar heads and hydrocarbon tails of the surfactant molecules (Schott, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 271).

5 The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions offer the advantage of
10 solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed spontaneously.

Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers, polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprates (MCA750), decaglycerol monooleate (MO750), decaglycerol sequioleate (SO750), decaglycerol decaoleate (DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into
20 the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-
25

-35-

emulsifying microemulsion systems are known in the art. The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives
5 of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain (C8-C12) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides,
10 saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone oil.

Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w
15 and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides (Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385-1390; Ritschel, *Meth. Find. Exp. Clin. Pharmacol.*, 1993, 13, 205). Microemulsions afford advantages of
20 improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms,
25 improved clinical potency, and decreased toxicity (Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385; Ho et al., *J. Pharm. Sci.*, 1996, 85, 138-143). Often microemulsions may form spontaneously when their components are brought together at ambient temperature. This may be
30 particularly advantageous when formulating thermolabile drugs, peptides or oligonucleotides. Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical

-36-

applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of oligonucleotides and nucleic acids from the gastrointestinal tract, as well as improve the local cellular uptake of oligonucleotides and nucleic acids within the gastrointestinal tract, vagina, buccal cavity and other areas of administration.

Microemulsions of the present invention may also contain additional components and additives such as sorbitan monostearate (Grill 3), Labrasol, and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the oligonucleotides and nucleic acids of the present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as belonging to one of five broad categories - surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p. 92). Each of these classes has been discussed above.

Liposomes

There are many organized surfactant structures besides microemulsions that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the duration of action they offer from the standpoint of drug delivery. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

-37-

Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess
5 the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as efficiently with the cell wall, are taken up by macrophages *in vivo*.

In order to cross intact mammalian skin, lipid
10 vesicles must pass through a series of fine pores, each with a diameter less than 50 nm, under the influence of a suitable transdermal gradient. Therefore, it is desirable to use a liposome which is highly deformable and able to pass through such fine pores.

15 Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from
20 metabolism and degradation (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size and
25 the aqueous volume of the liposomes.

Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the
30 liposomes start to merge with the cellular membranes. As the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

-38-

Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the skin. The majority of applications resulted in the targeting of the upper epidermis.

Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang et al., *Biochem. Biophys. Res. Commun.*, 1987, 147, 980-985).

Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous

-39-

gene was detected in the target cells (Zhou et al., *Journal of Controlled Release*, 1992, 19, 269-274).

One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

Several studies have assessed the topical delivery of liposomal drug formulations to the skin. Application of liposomes containing interferon to guinea pig skin resulted in a reduction of skin herpes sores while delivery of interferon via other means (e.g. as a solution or as an emulsion) were ineffective (Weiner et al., *Journal of Drug Targeting*, 1992, 2, 405-410). Further, an additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon using an aqueous system, and concluded that the liposomal formulation was superior to aqueous administration (du Plessis et al., *Antiviral Research*, 1992, 18, 259-265).

Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin, in particular systems comprising non-ionic surfactant and cholesterol. Non-ionic liposomal formulations comprising Novasome™ I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome™ II (glyceryl distearate/

-40-

cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the dermis of mouse skin.

Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of

5 cyclosporin-A into different layers of the skin (Hu et al. *S.T.P. Pharma. Sci.*, 1994, 4, 6, 466).

Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside G_{M1} , or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen et al., *FEBS Letters*, 1987, 223, 42; Wu et al., *Cancer Research*, 1993, 53, 3765). Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos et al. (*Ann. N.Y. Acad. Sci.*, 1987, 507, 64) reported the ability of monosialoganglioside G_{M1} , galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon et al. (*Proc. Natl. Acad. Sci. U.S.A.*, 1988, 85, 6949). U.S. Patent No. 4,837,028 and WO 88/04924, both to Allen et al., disclose liposomes

-41-

comprising (1) sphingomyelin and (2) the ganglioside G_{M1} or a galactocerebroside sulfate ester. U.S. Patent No.

5,543,152 (Webb et al.) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-sn-

5 dimyristoylphosphatidylcholine are disclosed in WO 97/13499 (Lim et al.).

Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto et al. (*Bull.*

10 *Chem. Soc. Jpn.*, 1980, 53, 2778) described liposomes comprising a nonionic detergent, 2C₁₂15G, that contains a PEG moiety. Illum et al. (*FEBS Lett.*, 1984, 167, 79) noted that hydrophilic coating of polystyrene particles with polymeric glycols results in significantly enhanced blood

15 half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Patent Nos.

4,426,330 and 4,534,899). Klivanov et al. (*FEBS Lett.*, 1990, 268, 235) described experiments demonstrating that

20 liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume et al. (*Biochimica et Biophysica Acta*, 1990, 1029, 91) extended such observations to other PEG-derivatized phospholipids,

25 e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG.

Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome

30 compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle et al. (U.S. Patent Nos. 5,013,556 and 5,356,633) and Martin et al. (U.S. Patent No. 5,213,804 and European

-42-

Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Patent No. 5,225,212 (both to Martin *et al.*) and in WO 94/20073 (Zalipsky *et al.*) Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi *et al.*). U.S. Patent Nos. 5,540,935 (Miyazaki *et al.*) and 5,556,948 (Tagawa *et al.*) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

10 A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry *et al.* discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Patent No. 5,264,221 to Tagawa *et al.* discloses protein-bonded liposomes and
15 asserts that the contents of such liposomes may include an antisense RNA. U.S. Patent No. 5,665,710 to Rahman *et al.* describes certain methods of encapsulating oligodeoxynucleotides in liposomes. WO 97/04787 to Love *et al.* discloses liposomes comprising antisense
20 oligonucleotides targeted to the raf gene.

Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly
25 deformable that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, *e.g.* they are self-optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their
30 targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum

-43-

albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

5 Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile
10 balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

15 If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values. In general their HLB values range from 2 to about 18 depending
20 on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates,
25 propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included in this class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

30 If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates

-44-

and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of this class.

If the surfactant molecule has the ability to carry either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

The use of surfactants in drug products, formulations and in emulsions has been reviewed (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

Penetration Enhancers

In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides, to the skin of animals. Most drugs are present in solution in both ionized and nonionized forms. However, usually only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to aiding the diffusion of non-lipophilic drugs

-45-

across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs.

Penetration enhancers may be classified as belonging to one of five broad categories, i.e., surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92). Each of the above mentioned classes of penetration enhancers are described below in greater detail.

10

Surfactants: In connection with the present invention, surfactants (or "surface-active agents") are chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid, with the result that absorption of oligonucleotides through the mucosa is enhanced. In addition to bile salts and fatty acids, these penetration enhancers include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92); and perfluorochemical emulsions, such as FC-43. Takahashi et al., *J. Pharm. Pharmacol.*, 1988, 40, 252).

25

Fatty acids: Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein (1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arachidonic acid, glycerol 1-monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines,

30

-46-

- C₁₋₁₀ alkyl esters thereof (e.g., methyl, isopropyl and t-butyl), and mono- and di-glycerides thereof (i.e., oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; El Hariri et al., *J. Pharm. Pharmacol.*, 1992, 44, 651-654).
- 10 Bile salts: The physiological role of bile includes the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman et al. Eds., McGraw-Hill, New York, 1996, pp. 15 934-935). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. The bile salts of the invention include, for 20 example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), glucolic acid (sodium glucolate), glycholic acid (sodium glycocholate), glycodeoxycholic acid (sodium 25 glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate), ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydro-fusidate (STDHF), sodium glycodihydrofusidate and 30 polyoxyethylene-9-lauryl ether (POE) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Swinyard, Chapter 39 In: *Remington's Pharmaceutical Sciences*, 18th Ed., Gennaro, ed., Mack Publishing Co.,

-47-

Easton, PA, 1990, pages 782-783; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Yamamoto et al., *J. Pharm. Exp. Ther.*, 1992, 263, 25; Yamashita et al., *J. Pharm. Sci.*, 1990, 79, 579-583).

5

Chelating Agents: Chelating agents, as used in connection with the present invention, can be defined as compounds that remove metallic ions from solution by forming complexes therewith, with the result that
10 absorption of oligonucleotides through the mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal
15 ion for catalysis and are thus inhibited by chelating agents (Jarrett, *J. Chromatogr.*, 1993, 618, 315-339). Chelating agents of the invention include but are not limited to disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-methoxysalicylate and homovanilate), *N*-acyl derivatives of
20 collagen, laureth-9 and *N*-amino acyl derivatives of beta-diketones (enamines) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990,
25 7, 1-33; Buur et al., *J. Control Rel.*, 1990, 14, 43-51).

Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant
30 activity as chelating agents or as surfactants but that nonetheless enhance absorption of oligonucleotides through the alimentary mucosa (Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33). This

-48-

class of penetration enhancers include, for example, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee *et al.*, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita *et al.*, *J. Pharm. Pharmacol.*, 1987, 39, 621-626).

Agents that enhance uptake of oligonucleotides at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi *et al.*, U.S. Patent No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo *et al.*, PCT Application WO 97/30731), are also known to enhance the cellular uptake of oligonucleotides.

Other agents may be utilized to enhance the penetration of the administered nucleic acids, including glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

Carriers

Certain compositions of the present invention also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a nucleic acid, or analog thereof, which is inert (*i.e.*, does not possess biological activity *per se*) but is recognized as a nucleic acid by *in vivo* processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance,

-49-

can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For example, the recovery of a partially phosphorothioate oligonucleotide in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'-isothiocyano-stilbene-2,2'-disulfonic acid (Miyao et al., *Antisense Res. Dev.*, 1995, 5, 115-121; Takakura et al., *Antisense & Nucl. Acid Drug Dev.*, 1996, 6, 177-183).

Excipients

In contrast to a carrier compound, a "pharmaceutical carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); lubricants (e.g., magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, etc.); disintegrants (e.g., starch, sodium starch

-50-

glycolate, etc.); and wetting agents (e.g., sodium lauryl sulphate, etc.).

Pharmaceutically acceptable organic or inorganic excipient suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous solutions, non-aqueous solutions in common solvents such as alcohols, or solutions of the nucleic acids in liquid or solid oil bases. The solutions may also contain buffers, diluents and other suitable additives. Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral administration which do not deleteriously react with nucleic acids can be used.

Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Other Components

The compositions of the present invention may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example,

-51-

antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as

5 dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations

10 can be sterilized and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact

15 with the nucleic acid(s) of the formulation.

Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

20 Certain embodiments of the invention provide pharmaceutical compositions containing (a) one or more antisense compounds and (b) one or more other chemotherapeutic agents which function by a non-antisense mechanism. Examples of such chemotherapeutic agents

25 include, but are not limited to, anticancer drugs such as daunorubicin, dactinomycin, doxorubicin, bleomycin, mitomycin, nitrogen mustard, chlorambucil, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine (CA), 5-fluorouracil (5-FU), floxuridine

30 (5-FUdR), methotrexate (MTX), colchicine, vincristine, vinblastine, etoposide, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds.,

-52-

1987, Rahway, N.J., pages 1206-1228). Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 2499-2506 and 46-49, respectively). Other non-antisense chemotherapeutic agents are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

In another related embodiment, compositions of the invention may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds targeted to a second nucleic acid target. Numerous examples of antisense compounds are known in the art. Two or more combined compounds may be used together or sequentially.

The formulation of therapeutic compositions and their subsequent administration is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can generally be estimated based on EC_{50} s found to be effective in *in vitro* and *in vivo* animal models. In general, dosage is from 0.01 ug to 100 g per kg of body

-53-

weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence
5 times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses,
10 ranging from 0.01 ug to 100 g per kg of body weight, once or more daily, to once every 20 years.

While the present invention has been described with specificity in accordance with certain of its preferred embodiments, the following examples serve only to
15 illustrate the invention and are not intended to limit the same.

EXAMPLES

20 Example 1

Nucleoside Phosphoramidites for Oligonucleotide Synthesis Deoxy and 2'-alkoxy amidites

2'-Deoxy and 2'-methoxy beta-cyanoethyl diisopropyl phosphoramidites were purchased from commercial sources
25 (e.g. Chemgenes, Needham MA or Glen Research, Inc. Sterling VA). Other 2'-O-alkoxy substituted nucleoside amidites are prepared as described in U.S. Patent 5,506,351, herein incorporated by reference. For oligonucleotides synthesized using 2'-alkoxy amidites, the standard cycle
30 for unmodified oligonucleotides was utilized, except the wait step after pulse delivery of tetrazole and base was increased to 360 seconds.

-54-

Oligonucleotides containing 5-methyl-2'-deoxycytidine (5-Me-C) nucleotides were synthesized according to published methods [Sanghvi, et. al., *Nucleic Acids Research*, 1993, 21, 3197-3203] using commercially available
5 phosphoramidites (Glen Research, Sterling VA or ChemGenes, Needham MA).

2'-Fluoro amidites

2'-Fluorodeoxyadenosine amidites

10 2'-fluoro oligonucleotides were synthesized as described previously [Kawasaki, et. al., *J. Med. Chem.*, 1993, 36, 831-841] and United States patent 5,670,633, herein incorporated by reference. Briefly, the protected nucleoside N6-benzoyl-2'-deoxy-2'-fluoroadenosine was
15 synthesized utilizing commercially available 9-beta-D-arabinofuranosyladenine as starting material and by modifying literature procedures whereby the 2'-alpha-fluoro atom is introduced by a S_N2-displacement of a 2'-beta-trityl group. Thus N6-benzoyl-9-beta-D-arabinofuranosyladenine
20 was selectively protected in moderate yield as the 3',5'-ditetrahydropyranyl (THP) intermediate. Deprotection of the THP and N6-benzoyl groups was accomplished using standard methodologies and standard methods were used to obtain the 5'-dimethoxytrityl-(DMT) and 5'-DMT-3'-
25 phosphoramidite intermediates.

2'-Fluorodeoxyguanosine

The synthesis of 2'-deoxy-2'-fluoroguanosine was accomplished using tetraisopropylidisiloxanyl (TPDS)
30 protected 9-beta-D-arabinofuranosylguanine as starting material, and conversion to the intermediate diisobutyryl-arabinofuranosylguanosine. Deprotection of the TPDS group was followed by protection of the hydroxyl group with THP

-55-

to give diisobutyryl di-THP protected arabinofuranosylguanine. Selective O-deacylation and triflation was followed by treatment of the crude product with fluoride, then deprotection of the THP groups.

5 Standard methodologies were used to obtain the 5'-DMT- and 5'-DMT-3'-phosphoramidites.

2'-Fluorouridine

Synthesis of 2'-deoxy-2'-fluorouridine was
10 accomplished by the modification of a literature procedure in which 2,2'-anhydro-1-beta-D-arabinofuranosyluracil was treated with 70% hydrogen fluoride-pyridine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

15

2'-Fluorodeoxycytidine

2'-deoxy-2'-fluorocytidine was synthesized via amination of 2'-deoxy-2'-fluorouridine, followed by selective protection to give N4-benzoyl-2'-deoxy-2'-
20 fluorocytidine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

2'-O-(2-Methoxyethyl) modified amidites

2'-O-Methoxyethyl-substituted nucleoside amidites are
25 prepared as follows, or alternatively, as per the methods of Martin, P., *Helvetica Chimica Acta*, 1995, 78, 486-504.

2,2'-Anhydro[1-(beta-D-arabinofuranosyl)-5-methyluridine]

30 5-Methyluridine (ribosylthymine, commercially available through Yamasa, Choshi, Japan) (72.0 g, 0.279 M), diphenylcarbonate (90.0 g, 0.420 M) and sodium bicarbonate (2.0 g, 0.024 M) were added to DMF (300 mL). The mixture

-56-

was heated to reflux, with stirring, allowing the evolved carbon dioxide gas to be released in a controlled manner. After 1 hour, the slightly darkened solution was concentrated under reduced pressure. The resulting syrup was poured into diethylether (2.5 L), with stirring. The product formed a gum. The ether was decanted and the residue was dissolved in a minimum amount of methanol (ca. 400 mL). The solution was poured into fresh ether (2.5 L) to yield a stiff gum. The ether was decanted and the gum was dried in a vacuum oven (60°C at 1 mm Hg for 24 h) to give a solid that was crushed to a light tan powder (57 g, 85% crude yield). The NMR spectrum was consistent with the structure, contaminated with phenol as its sodium salt (ca. 5%). The material was used as is for further reactions (or it can be purified further by column chromatography using a gradient of methanol in ethyl acetate (10-25%) to give a white solid, mp 222-4°C).

2'-O-Methoxyethyl-5-methyluridine

2,2'-Anhydro-5-methyluridine (195 g, 0.81 M), tris(2-methoxyethyl)borate (231 g, 0.98 M) and 2-methoxyethanol (1.2 L) were added to a 2 L stainless steel pressure vessel and placed in a pre-heated oil bath at 160°C. After heating for 48 hours at 155-160°C, the vessel was opened and the solution evaporated to dryness and triturated with MeOH (200 mL). The residue was suspended in hot acetone (1 L). The insoluble salts were filtered, washed with acetone (150 mL) and the filtrate evaporated. The residue (280 g) was dissolved in CH₃CN (600 mL) and evaporated. A silica gel column (3 kg) was packed in CH₂Cl₂/acetone/MeOH (20:5:3) containing 0.5% Et₃NH. The residue was dissolved in CH₂Cl₂ (250 mL) and adsorbed onto silica (150 g) prior to loading onto the column. The product was eluted with the packing

-57-

solvent to give 160 g (63%) of product. Additional material was obtained by reworking impure fractions.

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine

5 2'-O-Methoxyethyl-5-methyluridine (160 g, 0.506 M) was co-evaporated with pyridine (250 mL) and the dried residue dissolved in pyridine (1.3 L). A first aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the mixture stirred at room temperature for one hour. A
10 second aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the reaction stirred for an additional one hour. Methanol (170 mL) was then added to stop the reaction. HPLC showed the presence of approximately 70% product. The solvent was evaporated and triturated with
15 CH₃CN (200 mL). The residue was dissolved in CHCl₃ (1.5 L) and extracted with 2x500 mL of saturated NaHCO₃ and 2x500 mL of saturated NaCl. The organic phase was dried over Na₂SO₄, filtered and evaporated. 275 g of residue was obtained. The residue was purified on a 3.5 kg silica gel column,
20 packed and eluted with EtOAc/hexane/acetone (5:5:1) containing 0.5% Et₃NH. The pure fractions were evaporated to give 164 g of product. Approximately 20 g additional was obtained from the impure fractions to give a total yield of 183 g (57%).

25

3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (106 g, 0.167 M), DMF/pyridine (750 mL of a 3:1 mixture
30 prepared from 562 mL of DMF and 188 mL of pyridine) and acetic anhydride (24.38 mL, 0.258 M) were combined and stirred at room temperature for 24 hours. The reaction was monitored by TLC by first quenching the TLC sample with the

-58-

addition of MeOH. Upon completion of the reaction, as judged by TLC, MeOH (50 mL) was added and the mixture evaporated at 35°C. The residue was dissolved in CHCl₃ (800 mL) and extracted with 2x200 mL of saturated sodium bicarbonate and 2x200 mL of saturated NaCl. The water layers were back extracted with 200 mL of CHCl₃. The combined organics were dried with sodium sulfate and evaporated to give 122 g of residue (approx. 90% product).

The residue was purified on a 3.5 kg silica gel column and eluted using EtOAc/hexane(4:1). Pure product fractions were evaporated to yield 96 g (84%). An additional 1.5 g was recovered from later fractions.

15 3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine

A first solution was prepared by dissolving 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (96 g, 0.144 M) in CH₃CN (700 mL) and set aside. Triethylamine (189 mL, 1.44 M) was added to a solution of triazole (90 g, 1.3 M) in CH₃CN (1 L), cooled to -5°C and stirred for 0.5 h using an overhead stirrer. POCl₃ was added dropwise, over a 30 minute period, to the stirred solution maintained at 0-10°C, and the resulting mixture stirred for an additional 2 hours. The first solution was added dropwise, over a 45 minute period, to the latter solution. The resulting reaction mixture was stored overnight in a cold room. Salts were filtered from the reaction mixture and the solution was evaporated. The residue was dissolved in EtOAc (1 L) and the insoluble solids were removed by filtration. The filtrate was washed with 1x300 mL of NaHCO₃ and 2x300 mL of saturated NaCl, dried over sodium sulfate and evaporated. The residue was triturated with EtOAc to give the title compound.

-59-

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine

A solution of 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine (103 g, 0.141 M) in dioxane (500 mL) and NH_4OH (30 mL) was stirred at room temperature for 2 hours. The dioxane solution was evaporated and the residue azeotroped with MeOH (2x200 mL). The residue was dissolved in MeOH (300 mL) and transferred to a 2 liter stainless steel pressure vessel. MeOH (400 mL) saturated with NH_3 gas was added and the vessel heated to 100°C for 2 hours (TLC showed complete conversion). The vessel contents were evaporated to dryness and the residue was dissolved in EtOAc (500 mL) and washed once with saturated NaCl (200 mL). The organics were dried over sodium sulfate and the solvent was evaporated to give 85 g (95%) of the title compound.

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (85 g, 0.134 M) was dissolved in DMF (800 mL) and benzoic anhydride (37.2 g, 0.165 M) was added with stirring. After stirring for 3 hours, TLC showed the reaction to be approximately 95% complete. The solvent was evaporated and the residue azeotroped with MeOH (200 mL). The residue was dissolved in CHCl_3 (700 mL) and extracted with saturated NaHCO_3 (2x300 mL) and saturated NaCl (2x300 mL), dried over MgSO_4 and evaporated to give a residue (96 g). The residue was chromatographed on a 1.5 kg silica column using EtOAc/hexane (1:1) containing 0.5% Et_3NH as the eluting solvent. The pure product fractions were evaporated to give 90 g (90%) of the title compound.

-60-

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine-3'-amidite

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (74 g, 0.10 M) was dissolved in CH₂Cl₂ (1 L). Tetrazole diisopropylamine (7.1 g) and 2-cyanoethoxy-tetra-(isopropyl)phosphite (40.5 mL, 0.123 M) were added with stirring, under a nitrogen atmosphere. The resulting mixture was stirred for 20 hours at room temperature (TLC showed the reaction to be 95% complete). The reaction mixture was extracted with saturated NaHCO₃ (1x300 mL) and saturated NaCl (3x300 mL). The aqueous washes were back-extracted with CH₂Cl₂ (300 mL), and the extracts were combined, dried over MgSO₄ and concentrated. The residue obtained was chromatographed on a 1.5 kg silica column using EtOAc/hexane (3:1) as the eluting solvent. The pure fractions were combined to give 90.6 g (87%) of the title compound.

2'-O-(Aminooxyethyl) nucleoside amidites and 2'-O-(dimethylaminooxyethyl) nucleoside amidites

2'-(Dimethylaminooxyethoxy) nucleoside amidites

2'-(Dimethylaminooxyethoxy) nucleoside amidites [also known in the art as 2'-O-(dimethylaminooxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and guanosine nucleoside amidites are prepared similarly to the thymidine (5-methyluridine) except the exocyclic amines are protected with a benzoyl moiety in the case of adenosine and cytidine and with isobutyryl in the case of guanosine.

-61-

5'-O-tert-Butyldiphenylsilyl-O²-2'-anhydro-5-methyluridine

O²-2'-anhydro-5-methyluridine (Pro. Bio. Sint., Varese, Italy, 100.0g, 0.416 mmol), dimethylaminopyridine (0.66g, 0.013eq, 0.0054mmol) were dissolved in dry pyridine (500 ml) at ambient temperature under an argon atmosphere and with mechanical stirring. tert-Butyldiphenylchlorosilane (125.8g, 119.0mL, 1.1eq, 0.458mmol) was added in one portion. The reaction was stirred for 16 h at ambient temperature. TLC (Rf 0.22, ethyl acetate) indicated a complete reaction. The solution was concentrated under reduced pressure to a thick oil. This was partitioned between dichloromethane (1 L) and saturated sodium bicarbonate (2x1 L) and brine (1 L). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to a thick oil. The oil was dissolved in a 1:1 mixture of ethyl acetate and ethyl ether (600mL) and the solution was cooled to -10°C. The resulting crystalline product was collected by filtration, washed with ethyl ether (3x200 mL) and dried (40°C, 1mm Hg, 24 h) to 149g (74.8%) of white solid. TLC and NMR were consistent with pure product.

5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine

In a 2 L stainless steel, unstirred pressure reactor was added borane in tetrahydrofuran (1.0 M, 2.0 eq, 622 mL). In the fume hood and with manual stirring, ethylene glycol (350 mL, excess) was added cautiously at first until the evolution of hydrogen gas subsided. 5'-O-tert-Butyldiphenylsilyl-O²-2'-anhydro-5-methyluridine (149 g, 0.311 mol) and sodium bicarbonate (0.074 g, 0.003 eq) were added with manual stirring. The reactor was sealed and

-62-

added with manual stirring. The reactor was sealed and heated in an oil bath until an internal temperature of 160° C was reached and then maintained for 16 h (pressure < 100 psig). The reaction vessel was cooled to ambient and
5 opened. TLC (Rf 0.67 for desired product and Rf 0.82 for ara-T side product, ethyl acetate) indicated about 70% conversion to the product. In order to avoid additional side product formation, the reaction was stopped, concentrated under reduced pressure (10 to 1mm Hg) in a
10 warm water bath (40-100°C) with the more extreme conditions used to remove the ethylene glycol. [Alternatively, once the low boiling solvent is gone, the remaining solution can be partitioned between ethyl acetate and water. The product will be in the organic phase.] The residue was
15 purified by column chromatography (2kg silica gel, ethyl acetate-hexanes gradient 1:1 to 4:1). The appropriate fractions were combined, stripped and dried to product as a white crisp foam (84g, 50%), contaminated starting material (17.4g) and pure reusable starting material 20g. The yield
20 based on starting material less pure recovered starting material was 58%. TLC and NMR were consistent with 99% pure product.

2'-O-([2-phthalimidoxy)ethyl]-5'-t-butylidiphenylsilyl-5-methyluridine

25 5'-O-tert-Butylidiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine (20g, 36.98mmol) was mixed with triphenylphosphine (11.63g, 44.36mmol) and N-hydroxyphthalimide (7.24g, 44.36mmol). It was then dried over P₂O₅ under high vacuum for two days at 40°C. The
30 reaction mixture was flushed with argon and dry THF (369.8mL, Aldrich, sure seal bottle) was added to get a clear solution. Diethyl-azodicarboxylate (6.98mL, 44.36mmol) was added dropwise to the reaction mixture. The

-63-

rate of addition is maintained such that resulting deep red coloration is just discharged before adding the next drop. After the addition was complete, the reaction was stirred for 4 hrs. By that time TLC showed the completion of the reaction (ethylacetate:hexane, 60:40). The solvent was evaporated in vacuum. Residue obtained was placed on a flash column and eluted with ethyl acetate:hexane (60:40), to get 2'-O-[(2-phthalimidooxy)ethyl]-5'-t-butyl-
5 10 86%).

5'-O-tert-butyl-diphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine

2'-O-[(2-phthalimidooxy)ethyl]-5'-t-butyl-diphenylsilyl-5-methyluridine (3.1g, 4.5mmol) was dissolved in dry CH₂Cl₂ (4.5mL) and methylhydrazine (300mL, 4.64mmol) was added dropwise at -10°C to 0°C. After 1 h the mixture was filtered, the filtrate was washed with ice cold CH₂Cl₂ and the combined organic phase was washed with water, brine and dried over anhydrous Na₂SO₄. The solution was concentrated to get 2'-O-(aminooxyethyl) thymidine, which was then dissolved in MeOH (67.5mL). To this formaldehyde (20% aqueous solution, w/w, 1.1 eq.) was added and the resulting mixture was stirred for 1 h. Solvent was removed under vacuum; residue chromatographed to get 5'-O-tert-butyl-diphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine as white foam (1.95 g, 78%).

5'-O-tert-Butyl-diphenylsilyl-2'-O-[N,N-dimethylaminooxyethyl]-5-methyluridine

5'-O-tert-butyl-diphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine (1.77g, 3.12mmol) was dissolved in a solution of 1M pyridinium p-

-64-

toluenesulfonate (PPTS) in dry MeOH (30.6mL). Sodium cyanoborohydride (0.39g, 6.13mmol) was added to this solution at 10°C under inert atmosphere. The reaction mixture was stirred for 10 minutes at 10°C. After that the reaction vessel was removed from the ice bath and stirred at room temperature for 2 h, the reaction monitored by TLC (5% MeOH in CH₂Cl₂). Aqueous NaHCO₃ solution (5%, 10mL) was added and extracted with ethyl acetate (2x20mL). Ethyl acetate phase was dried over anhydrous Na₂SO₄, evaporated to dryness. Residue was dissolved in a solution of 1M PPTS in MeOH (30.6mL). Formaldehyde (20% w/w, 30mL, 3.37mmol) was added and the reaction mixture was stirred at room temperature for 10 minutes. Reaction mixture cooled to 10°C in an ice bath, sodium cyanoborohydride (0.39g, 6.13mmol) was added and reaction mixture stirred at 10°C for 10 minutes. After 10 minutes, the reaction mixture was removed from the ice bath and stirred at room temperature for 2 hrs. To the reaction mixture 5% NaHCO₃ (25mL) solution was added and extracted with ethyl acetate (2x25mL). Ethyl acetate layer was dried over anhydrous Na₂SO₄ and evaporated to dryness. The residue obtained was purified by flash column chromatography and eluted with 5% MeOH in CH₂Cl₂ to get 5'-O-tert-butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine as a white foam (14.6g, 80%).

2'-O-(dimethylaminoxyethyl)-5-methyluridine

Triethylamine trihydrofluoride (3.91mL, 24.0mmol) was dissolved in dry THF and triethylamine (1.67mL, 12mmol, dry, kept over KOH). This mixture of triethylamine-2HF was then added to 5'-O-tert-butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine (1.40g, 2.4mmol) and stirred at room temperature for 24 hrs. Reaction was

-65-

monitored by TLC (5% MeOH in CH₂Cl₂). Solvent was removed under vacuum and the residue placed on a flash column and eluted with 10% MeOH in CH₂Cl₂ to get 2'-O-(dimethylaminooxyethyl)-5-methyluridine (766mg, 92.5%).

5

5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine 2'-O-(dimethylaminoxyethyl)-5-methyluridine (750mg, 2.17mmol) was dried over P₂O₅ under high vacuum overnight at 40°C. It was then co-evaporated with anhydrous pyridine (20mL). The residue obtained was dissolved in pyridine (11mL) under argon atmosphere. 4-dimethylaminopyridine (26.5mg, 2.60mmol), 4,4'-dimethoxytrityl chloride (880mg, 2.60mmol) was added to the mixture and the reaction mixture was stirred at room temperature until all of the starting material disappeared. Pyridine was removed under vacuum and the residue chromatographed and eluted with 10% MeOH in CH₂Cl₂ (containing a few drops of pyridine) to get 5'-O-DMT-2'-O-(dimethylamino-oxyethyl)-5-methyluridine (1.13g, 80%).

20

5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]
5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine (1.08g, 1.67mmol) was co-evaporated with toluene (20mL). To the residue N,N-diisopropylamine tetrazonide (0.29g, 1.67mmol) was added and dried over P₂O₅ under high vacuum overnight at 40°C. Then the reaction mixture was dissolved in anhydrous acetonitrile (8.4mL) and 2-cyanoethyl-N,N,N¹,N¹-tetraisopropylphosphoramidite (2.12mL, 6.08mmol) was added. The reaction mixture was stirred at ambient temperature for 4 hrs under inert atmosphere. The progress of the reaction was monitored by TLC (hexane:ethyl acetate

30

-66-

1:1). The solvent was evaporated, then the residue was dissolved in ethyl acetate (70mL) and washed with 5% aqueous NaHCO₃ (40mL). Ethyl acetate layer was dried over anhydrous Na₂SO₄ and concentrated. Residue obtained was chromatographed (ethyl acetate as eluent) to get 5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite] as a foam (1.04g, 74.9%).

10 2'-(Aminooxyethoxy) nucleoside amidites

2'-(Aminooxyethoxy) nucleoside amidites [also known in the art as 2'-O-(aminooxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and thymidine nucleoside amidites are prepared similarly.

N2-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]

20 The 2'-O-aminooxyethyl guanosine analog may be obtained by selective 2'-O-alkylation of diaminopurine riboside. Multigram quantities of diaminopurine riboside may be purchased from Schering AG (Berlin) to provide 2'-O-(2-ethylacetyl) diaminopurine riboside along with a minor amount of the 3'-O-isomer. 2'-O-(2-ethylacetyl) diaminopurine riboside may be resolved and converted to 2'-O-(2-ethylacetyl)guanosine by treatment with adenosine deaminase. (McGee, D. P. C., Cook, P. D., Guinasso, C. J., WO 94/02501 A1 940203.) Standard protection procedures should afford 2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine and 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine which may be reduced to provide

-67-

2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-
5'-O-(4,4'-dimethoxytrityl)guanosine. As before the
hydroxyl group may be displaced by N-hydroxyphthalimide via
a Mitsunobu reaction, and the protected nucleoside may
5 phosphitylated as usual to yield 2-N-isobutyryl-6-O-
diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-
dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-
diisopropylphosphoramidite].

10 2'-dimethylaminoethoxyethoxy (2'-DMAEOE) nucleoside
amidites

2'-dimethylaminoethoxyethoxy nucleoside amidites (also
known in the art as 2'-O-dimethylaminoethoxyethyl, i.e.,
2'-O-CH₂-O-CH₂-N(CH₂)₂, or 2'-DMAEOE nucleoside amidites) are
15 prepared as follows. Other nucleoside amidites are prepared
similarly.

2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine

2[2-(Dimethylamino)ethoxy]ethanol (Aldrich, 6.66 g, 50
20 mmol) is slowly added to a solution of borane in tetra-
hydrofuran (1 M, 10 mL, 10 mmol) with stirring in a 100 mL
bomb. Hydrogen gas evolves as the solid dissolves. O²-,2'-
anhydro-5-methyluridine (1.2 g, 5 mmol), and sodium
bicarbonate (2.5 mg) are added and the bomb is sealed,
25 placed in an oil bath and heated to 155°C for 26 hours. The
bomb is cooled to room temperature and opened. The crude
solution is concentrated and the residue partitioned
between water (200 mL) and hexanes (200 mL). The excess
phenol is extracted into the hexane layer. The aqueous
30 layer is extracted with ethyl acetate (3x200 mL) and the
combined organic layers are washed once with water, dried
over anhydrous sodium sulfate and concentrated. The
residue is columned on silica gel using methanol/methylene

-68-

chloride 1:20 (which has 2% triethylamine) as the eluent. As the column fractions are concentrated a colorless solid forms which is collected to give the title compound as a white solid.

5

5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]]-5-methyl uridine

To 0.5 g (1.3 mmol) of 2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]]-5-methyl uridine in anhydrous pyridine (8 mL), triethylamine (0.36 mL) and dimethoxytrityl chloride (DMT-Cl, 0.87 g, 2 eq.) are added and stirred for 1 hour. The reaction mixture is poured into water (200 mL) and extracted with CH₂Cl₂ (2x200 mL). The combined CH₂Cl₂ layers are washed with saturated NaHCO₃ solution, followed by saturated NaCl solution and dried over anhydrous sodium sulfate. Evaporation of the solvent followed by silica gel chromatography using MeOH:CH₂Cl₂:Et₃N (20:1, v/v, with 1% triethylamine) gives the title compound.

20 **5'-O-Dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]]-5-methyl uridine-3'-O-(cyanoethyl-N,N-diisopropyl)phosphoramidite**

Diisopropylaminotetrazolide (0.6 g) and 2-cyanoethoxy-N,N-diisopropyl phosphoramidite (1.1 mL, 2 eq.) are added to a solution of 5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]]-5-methyluridine (2.17 g, 3 mmol) dissolved in CH₂Cl₂ (20 mL) under an atmosphere of argon. The reaction mixture is stirred overnight and the solvent evaporated. The resulting residue is purified by silica gel flash column chromatography with ethyl acetate as the eluent to give the title compound.

30

-69-

Example 2**Oligonucleotide synthesis**

Unsubstituted and substituted phosphodiester (P=O) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard
5 phosphoramidite chemistry with oxidation by iodine.

Phosphorothioates (P=S) are synthesized as for the phosphodiester oligonucleotides except the standard oxidation bottle was replaced by 0.2 M solution of 3H-1,2-
10 benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation wait step was increased to 68 sec and was followed by the capping step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at 55°C (18
15 h), the oligonucleotides were purified by precipitating twice with 2.5 volumes of ethanol from a 0.5 M NaCl solution. Phosphinate oligonucleotides are prepared as described in U.S. Patent 5,508,270, herein incorporated by reference.

20 Alkyl phosphonate oligonucleotides are prepared as described in U.S. Patent 4,469,863, herein incorporated by reference.

3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Patents 5,610,289 or
25 5,625,050, herein incorporated by reference.

Phosphoramidite oligonucleotides are prepared as described in U.S. Patent, 5,256,775 or U.S. Patent 5,366,878, herein incorporated by reference.

Alkylphosphonothioate oligonucleotides are prepared as
30 described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively), herein incorporated by reference.

-70-

3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Patent 5,476,925, herein incorporated by reference.

Phosphotriester oligonucleotides are prepared as described in U.S. Patent 5,023,243, herein incorporated by reference.

Borano phosphate oligonucleotides are prepared as described in U.S. Patents 5,130,302 and 5,177,198, both herein incorporated by reference.

10

Example 3

Oligonucleoside Synthesis

Methylenemethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedi-
15 methylhydrazo linked oligonucleosides, also identified as MDH linked oligonucleosides, and methylenecarbonylamino linked oligonucleosides, also identified as amide-3 linked oligonucleosides, and methyleneaminocarbonyl linked oligonucleosides, also identified as amide-4 linked oligonucleo-
20 sides, as well as mixed backbone compounds having, for instance, alternating MMI and P=O or P=S linkages are prepared as described in U.S. Patents 5,378,825, 5,386,023, 5,489,677, 5,602,240 and 5,610,289, all of which are herein incorporated by reference.

25 Formacetal and thioformacetal linked oligonucleosides are prepared as described in U.S. Patents 5,264,562 and 5,264,564, herein incorporated by reference.

Ethylene oxide linked oligonucleosides are prepared as described in U.S. Patent 5,223,618, herein incorporated by
30 reference.

-71-

Example 4**PNA Synthesis**

Peptide nucleic acids (PNAs) are prepared in accordance with any of the various procedures referred to in Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications, *Bioorganic & Medicinal Chemistry*, 1996, 4, 5-23. They may also be prepared in accordance with U.S. Patents 5,539,082, 5,700,922, and 5,719,262, herein incorporated by reference.

10

Example 5**Synthesis of Chimeric Oligonucleotides**

Chimeric oligonucleotides, oligonucleosides or mixed oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3' "wing" segments of linked nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are also known in the art as "hemimers" or "wingmers".

25 [2'-O-Me]--[2'-deoxy]--[2'-O-Me] Chimeric**Phosphorothioate Oligonucleotides**

Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligonucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 380B, as above. Oligonucleotides are synthesized using the automated synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphoramidite for the DNA portion and 5'-dimethoxytrityl-2'-O-

30

-72-

5 methyl-3'-O-phosphoramidite for 5' and 3' wings. The standard synthesis cycle is modified by increasing the wait step after the delivery of tetrazole and base to 600 s repeated four times for RNA and twice for 2'-O-methyl. The fully protected oligonucleotide is cleaved from the support and the phosphate group is deprotected in 3:1 ammonia/ethanol at room temperature overnight then lyophilized to dryness. Treatment in methanolic ammonia for 24 hrs at room temperature is then done to deprotect all bases and sample was again lyophilized to dryness. The pellet is resuspended in 1M TBAF in THF for 24 hrs at room temperature to deprotect the 2' positions. The reaction is then quenched with 1M TEAA and the sample is then reduced to 1/2 volume by rotovac before being desalted on a G25 size exclusion column. The oligo recovered is then analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry.

20 [2'-O-(2-Methoxyethyl)]--[2'-deoxy]--[2'-O-(Methoxyethyl)] Chimeric Phosphorothioate Oligonucleotides

[2'-O-(2-methoxyethyl)]--[2'-deoxy]--[2'-O-(methoxyethyl)] chimeric phosphorothioate oligonucleotides were prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites.

30 [2'-O-(2-Methoxyethyl)Phosphodiester]--[2'-deoxy Phosphorothioate]--[2'-O-(2-Methoxyethyl) Phosphodiester] Chimeric Oligonucleotides

[2'-O-(2-methoxyethyl phosphodiester)]--[2'-deoxy phosphorothioate]--[2'-O-(methoxyethyl) phosphodiester] chimeric oligonucleotides are prepared as per the above

-73-

procedure for the 2'-O-methyl chimeric oligonucleotide with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites, oxidization with iodine to generate the phosphodiester internucleotide linkages within the wing portions of the chimeric structures and sulfurization utilizing 3,4-dihydro-2H-benzothiole-3-one 1,1-dioxide (Beaucage Reagent) to generate the phosphorothioate internucleotide linkages for the center gap.

Other chimeric oligonucleotides, chimeric oligonucleosides and mixed chimeric oligonucleotides/oligonucleosides are synthesized according to United States patent 5,623,065, herein incorporated by reference.

Example 6

15 Oligonucleotide Isolation

After cleavage from the controlled pore glass column (Applied Biosystems) and deblocking in concentrated ammonium hydroxide at 55°C for 18 hours, the oligonucleotides or oligonucleosides are purified by precipitation twice out of 0.5 M NaCl with 2.5 volumes ethanol. Synthesized oligonucleotides were analyzed by polyacrylamide gel electrophoresis on denaturing gels and judged to be at least 85% full length material. The relative amounts of phosphorothioate and phosphodiester linkages obtained in synthesis were periodically checked by ³¹P nuclear magnetic resonance spectroscopy, and for some studies oligonucleotides were purified by HPLC, as described by Chiang et al., *J. Biol. Chem.* 1991, 266, 18162-18171. Results obtained with HPLC-purified material were similar to those obtained with non-HPLC purified material.

-74-

Example 7**Oligonucleotide Synthesis - 96 Well Plate Format**

Oligonucleotides were synthesized via solid phase P(III) phosphoramidite chemistry on an automated synthesizer capable of assembling 96 sequences simultaneously in a standard 96 well format. Phosphodiester internucleotide linkages were afforded by oxidation with aqueous iodine. Phosphorothioate internucleotide linkages were generated by sulfurization utilizing 3,4-dihydro-2H-benzothiole-3-one 1,1-dioxide (Beaucage Reagent) in anhydrous acetonitrile. Standard base-protected beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial vendors (e.g. PE-Applied Biosystems, Foster City, CA, or Pharmacia, Piscataway, NJ). Non-standard nucleosides are synthesized as per known literature or patented methods. They are utilized as base-protected beta-cyanoethyl-diisopropyl phosphoramidites.

Oligonucleotides were cleaved from support and deprotected with concentrated NH_4OH at elevated temperature (55-60°C) for 12-16 hours and the released product then dried in vacuo. The dried product was then re-suspended in sterile water to afford a master plate from which all analytical and test plate samples are then diluted utilizing robotic pipettors.

25

Example 8**Oligonucleotide Analysis - 96 Well Plate Format**

The concentration of oligonucleotide in each well was assessed by dilution of samples and UV absorption spectroscopy. The full-length integrity of the individual products was evaluated by capillary electrophoresis (CE) in either the 96 well format (Beckman P/ACE™ MDQ) or, for individually prepared samples, on a commercial CE apparatus

30

-75-

(e.g., Beckman P/ACE™ 5000, ABI 270). Base and backbone composition was confirmed by mass analysis of the compounds utilizing electrospray-mass spectroscopy. All assay test plates were diluted from the master plate using single and multi-channel robotic pipettors. Plates were judged to be acceptable if at least 85% of the compounds on the plate were at least 85% full length.

Example 9

10 Cell culture and oligonucleotide treatment

The effect of antisense compounds on target nucleic acid expression can be tested in any of a variety of cell types provided that the target nucleic acid is present at measurable levels. This can be routinely determined using, for example, PCR or Northern blot analysis. The following 5 cell types are provided for illustrative purposes, but other cell types can be routinely used, provided that the target is expressed in the cell type chosen. This can be readily determined by methods routine in the art, for example Northern blot analysis, Ribonuclease protection assays, or RT-PCR.

T-24 cells:

The human transitional cell bladder carcinoma cell line T-24 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). T-24 cells were routinely cultured in complete McCoy's 5A basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached

-76-

90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analysis, cells may be
5 seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

A549 cells:

10 The human lung carcinoma cell line A549 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). A549 cells were routinely cultured in DMEM basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life
15 Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence.

20

NHDF cells:

Human neonatal dermal fibroblast (NHDF) were obtained from the Clonetics Corporation (Walkersville MD). NHDFs were routinely maintained in Fibroblast Growth Medium
25 (Clonetics Corporation, Walkersville MD) supplemented as recommended by the supplier. Cells were maintained for up to 10 passages as recommended by the supplier.

HEK cells:

30 Human embryonic keratinocytes (HEK) were obtained from the Clonetics Corporation (Walkersville MD). HEKs were routinely maintained in Keratinocyte Growth Medium (Clonetics Corporation, Walkersville MD) formulated as

-77-

recommended by the supplier. Cells were routinely maintained for up to 10 passages as recommended by the supplier.

5 MCF-7 cells:

The human breast carcinoma cell line MCF-7 was obtained from the American Type Culture Collection (Manassas, VA). MCF-7 cells were routinely cultured in DMEM low glucose (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analyses, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

20 Treatment with antisense compounds:

When cells reached 80% confluency, they were treated with oligonucleotide. For cells grown in 96-well plates, wells were washed once with 200 μ L OPTI-MEMTM-1 reduced-serum medium (Gibco BRL) and then treated with 130 μ L of OPTI-MEMTM-1 containing 3.75 μ g/mL LIPOFECTINTM (Gibco BRL) and the desired concentration of oligonucleotide. After 4-7 hours of treatment, the medium was replaced with fresh medium. Cells were harvested 16-24 hours after oligonucleotide treatment.

The concentration of oligonucleotide used varies from cell line to cell line. To determine the optimal oligonucleotide concentration for a particular cell line,

-78-

the cells are treated with a positive control oligonucleotide at a range of concentrations. For human cells the positive control oligonucleotide is ISIS 13920, TCCGTCATCGCTCCTCAGGG, SEQ ID NO: 1, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone which is targeted to human H-ras.

For mouse or rat cells the positive control oligonucleotide is ISIS 15770, ATGCATTCTGCCCCCAAGGA, SEQ ID NO: 2, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone which is targeted to both mouse and rat c-raf. The concentration of positive control oligonucleotide that results in 80% inhibition of c-Ha-ras (for ISIS 13920) or c-raf (for ISIS 15770) mRNA is then utilized as the screening concentration for new oligonucleotides in subsequent experiments for that cell line. If 80% inhibition is not achieved, the lowest concentration of positive control oligonucleotide that results in 60% inhibition of H-ras or c-raf mRNA is then utilized as the oligonucleotide screening concentration in subsequent experiments for that cell line. If 60% inhibition is not achieved, that particular cell line is deemed as unsuitable for oligonucleotide transfection experiments.

Example 10

Analysis of oligonucleotide inhibition of Her-2 expression

Antisense modulation of Her-2 expression can be assayed in a variety of ways known in the art. For example, Her-2 mRNA levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR). Real-time quantitative PCR is presently preferred. RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA.

-79-

Methods of RNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Northern blot analysis is

5 routine in the art and is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.2.1-4.2.9, John Wiley & Sons, Inc., 1996. Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7700 Sequence

10 Detection System, available from PE-Applied Biosystems, Foster City, CA and used according to manufacturer's instructions. Prior to quantitative PCR analysis, primer-probe sets specific to the target gene being measured are evaluated for their ability to be "multiplexed" with a

15 GAPDH amplification reaction. In multiplexing, both the target gene and the internal standard gene GAPDH are amplified concurrently in a single sample. In this analysis, mRNA isolated from untreated cells is serially diluted. Each dilution is amplified in the presence of

20 primer-probe sets specific for GAPDH only, target gene only ("single-plexing"), or both (multiplexing). Following PCR amplification, standard curves of GAPDH and target mRNA signal as a function of dilution are generated from both the single-plexed and multiplexed samples. If both the

25 slope and correlation coefficient of the GAPDH and target signals generated from the multiplexed samples fall within 10% of their corresponding values generated from the single-plexed samples, the primer-probe set specific for that target is deemed as multiplexable. Other methods of

30 PCR are also known in the art.

Protein levels of Her-2 can be quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis

-80-

(immunoblotting), ELISA or fluorescence-activated cell sorting (FACS). Antibodies directed to Her-2 can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies (Aerie Corporation, Birmingham, MI), or can be prepared via conventional antibody generation methods. Methods for preparation of polyclonal antisera are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.12.1-11.12.9, John Wiley & Sons, Inc., 1997.

Preparation of monoclonal antibodies is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.4.1-11.11.5, John Wiley & Sons, Inc., 1997.

Immunoprecipitation methods are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.16.1-10.16.11, John Wiley & Sons, Inc., 1998. Western blot (immunoblot) analysis is standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.8.1-10.8.21, John Wiley & Sons, Inc., 1997. Enzyme-linked immunosorbent assays (ELISA) are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.2.1-11.2.22, John Wiley & Sons, Inc., 1991.

Example 11

Poly(A)+ mRNA isolation

Poly(A)+ mRNA was isolated according to Miura et al., *Clin. Chem.*, 1996, 42, 1758-1764. Other methods for poly(A)+ mRNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.5.1-4.5.3, John Wiley & Sons,

-81-

Inc., 1993. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200 μ L cold PBS. 60 μ L lysis buffer (10 mM Tris-HCl, pH 7.6, 1 mM EDTA, 0.5 M NaCl, 0.5% NP-40, 20 mM
5 vanadyl-ribonucleoside complex) was added to each well, the plate was gently agitated and then incubated at room temperature for five minutes. 55 μ L of lysate was transferred to Oligo d(T) coated 96-well plates (AGCT Inc., Irvine CA). Plates were incubated for 60 minutes at room
10 temperature, washed 3 times with 200 μ L of wash buffer (10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl). After the final wash, the plate was blotted on paper towels to remove excess wash buffer and then air-dried for 5 minutes. 60 μ L of elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70°C
15 was added to each well, the plate was incubated on a 90°C hot plate for 5 minutes, and the eluate was then transferred to a fresh 96-well plate.

Cells grown on 100 mm or other standard plates may be treated similarly, using appropriate volumes of all
20 solutions.

Example 12

Total RNA Isolation

Total mRNA was isolated using an RNEASY 96™ kit and
25 buffers purchased from Qiagen Inc. (Valencia CA) following the manufacturer's recommended procedures. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200 μ L cold PBS. 100 μ L Buffer RLT was added to each well and the plate
30 vigorously agitated for 20 seconds. 100 μ L of 70% ethanol was then added to each well and the contents mixed by pipetting three times up and down. The samples were then

-82-

transferred to the RNEASY 96™ well plate attached to a QIAVAC™ manifold fitted with a waste collection tray and attached to a vacuum source. Vacuum was applied for 15 seconds. 1 mL of Buffer RW1 was added to each well of the RNEASY 96™ plate and the vacuum again applied for 15 seconds. 1 mL of Buffer RPE was then added to each well of the RNEASY 96™ plate and the vacuum applied for a period of 15 seconds. The Buffer RPE wash was then repeated and the vacuum was applied for an additional 10 minutes. The plate was then removed from the QIAVAC™ manifold and blotted dry on paper towels. The plate was then re-attached to the QIAVAC™ manifold fitted with a collection tube rack containing 1.2 mL collection tubes. RNA was then eluted by pipetting 60 µL water into each well, incubating 1 minute, and then applying the vacuum for 30 seconds. The elution step was repeated with an additional 60 µL water.

The repetitive pipetting and elution steps may be automated using a QIAGEN Bio-Robot 9604 (Qiagen, Inc., Valencia CA). Essentially, after lysing of the cells on the culture plate, the plate is transferred to the robot deck where the pipetting, DNase treatment and elution steps are carried out.

Example 13

Real-time Quantitative PCR Analysis of Her-2 mRNA Levels

Quantitation of Her-2 mRNA levels was determined by real-time quantitative PCR using the ABI PRISM™ 7700 Sequence Detection System (PE-Applied Biosystems, Foster City, CA) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of

-83-

polymerase chain reaction (PCR) products in real-time. As opposed to standard PCR, in which amplification products are quantitated after the PCR is completed, products in real-time quantitative PCR are quantitated as they

5 accumulate. This is accomplished by including in the PCR reaction an oligonucleotide probe that anneals specifically between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye (e.g., JOE, FAM, or VIC, obtained from either Operon Technologies Inc.,

10 Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 5' end of the probe and a quencher dye (e.g., TAMRA, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 3' end of the probe. When the probe

15 and dyes are intact, reporter dye emission is quenched by the proximity of the 3' quencher dye. During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-exonuclease activity of Taq polymerase. During the

20 extension phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated. With each cycle, additional reporter dye

25 molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular intervals by laser optics built into the ABI PRISM™ 7700 Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from

30 untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

-84-

PCR reagents were obtained from PE-Applied Biosystems, Foster City, CA. RT-PCR reactions were carried out by adding 25 μ L PCR cocktail (1x TAQMANTM buffer A, 5.5 mM MgCl₂, 300 μ M each of dATP, dCTP and dGTP, 600 μ M of dUTP, 5 100 nM each of forward primer, reverse primer, and probe, 20 Units RNase inhibitor, 1.25 Units AMPLITAQ GOLDTM, and 12.5 Units MuLV reverse transcriptase) to 96 well plates containing 25 μ L poly(A) mRNA solution. The RT reaction was carried out by incubation for 30 minutes at 48°C.

10 Following a 10 minute incubation at 95°C to activate the AMPLITAQ GOLDTM, 40 cycles of a two-step PCR protocol were carried out: 95°C for 15 seconds (denaturation) followed by 60°C for 1.5 minutes (annealing/extension).

Probes and primers to human Her-2 were designed to 15 hybridize to a human Her-2 sequence, using published sequence information (GenBank accession number X03363, incorporated herein as SEQ ID NO:3). For human Her-2 the PCR primers were:

forward primer: GTGACAGCAGAGGATGGAACAC (SEQ ID NO: 4)

20 reverse primer: ATGCCAGACCATAGCACACT (SEQ ID NO: 5) and the PCR probe was: FAM-CGGTGTGAGAAGTGCAGCAAGCCC-TAMRA (SEQ ID NO: 6) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

25 For human GAPDH the PCR primers were:

forward primer: CAACGGATTTGGTCGTATTGG (SEQ ID NO: 7)

reverse primer: GGCAACAATATCCACTTTACCAGAGT (SEQ ID NO: 8)

and the PCR probe was: 5' JOE-CGCCTGGTCACCAGGGCTGCT- TAMRA 3' (SEQ ID NO: 9) where JOE (PE-Applied Biosystems, Foster 30 City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

-85-

Example 14**Northern blot analysis of Her-2 mRNA levels**

Eighteen hours after antisense treatment, cell
5 monolayers were washed twice with cold PBS and lysed in 1
mL RNAzol™ (TEL-TEST "B" Inc., Friendswood, TX). Total RNA
was prepared following manufacturer's recommended
protocols. Twenty micrograms of total RNA was fractionated
by electrophoresis through 1.2% agarose gels containing
10 1.1% formaldehyde using a MOPS buffer system (AMRESCO, Inc.
Solon, OH). RNA was transferred from the gel to HYBOND™-N+
nylon membranes (Amersham Pharmacia Biotech, Piscataway,
NJ) by overnight capillary transfer using a
Northern/Southern Transfer buffer system (TEL-TEST "B"
15 Inc., Friendswood, TX). RNA transfer was confirmed by UV
visualization. Membranes were fixed by UV cross-linking
using a STRATALINKER™ UV Crosslinker 2400 (Stratagene,
Inc, La Jolla, CA) and then probed using QUICKHYB™
hybridization solution (Stratagene, La Jolla, CA) using
20 manufacturer's recommendations for stringent conditions.

To detect human Her-2, a human Her-2 specific probe
was prepared by PCR using the forward primer
GTGACAGCAGAGGATGGAACAC (SEQ ID NO: 4) and the reverse
primer ATGCCAGACCATAGCACACT (SEQ ID NO: 5). To normalize
25 for variations in loading and transfer efficiency membranes
were stripped and probed for human glyceraldehyde-3-
phosphate dehydrogenase (GAPDH) RNA (Clontech, Palo Alto,
CA).

Hybridized membranes were visualized and quantitated
30 using a PHOSPHORIMAGER™ and IMAGEQUANT™ Software V3.3
(Molecular Dynamics, Sunnyvale, CA). Data was normalized
to GAPDH levels in untreated controls.

-86-

Example 15

Antisense inhibition of human Her-2 expression by phosphorothioate oligodeoxynucleotides

5 In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human Her-2 RNA, using published sequences (GenBank accession number X03363, incorporated herein as SEQ ID NO: 3). The oligonucleotides are shown in Table 1. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 1 are are oligodeoxynucleotides with phosphorothioate backbones (internucleoside linkages) throughout. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on human Her-2 mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments. If present, "N.D." indicates "no data".

20

Table 1

Inhibition of human Her-2 mRNA levels by phosphorothioate oligodeoxynucleotides

ISIS #	REGION	TARGET (SEQ ID NO)	TARGET SITE	SEQUENCE	%INHIB	SEQ ID NO
27898	Coding	3	280	ggtctcgggactggcagg	32	10
27899	Coding	3	310	ctggtagagggtggcggag	0	11
27900	Coding	3	326	gcaccacctggcagccct	27	12
27901	Coding	3	365	tggcattggtgggcaggt	3	13
27902	Coding	3	379	caggaaggacaggctggc	6	14
27903	Coding	3	437	cctgcctcacttggttgt	36	15
27904	Coding	3	501	gccagggcatagttgtcc	19	16
27905	Coding	3	539	gggtggtattgttcagcg	6	17
27906	Coding	3	692	gctggttggtcttgtgga	27	18
27907	Coding	3	769	gcagcgggagcccttaca	17	19
27908	Coding	3	937	gtggttgaagtggaggca	2	20
27909	Coding	3	1159	gcacttctcacaccgctg	49	21

-87-

27910	Coding	3	1190	gaccatagcacactcggg	43	22
27911	Coding	3	1232	cactggtaactgccctca	46	23
27912	Coding	3	1327	ggcagtgttgaggctgg	43	24
27913	Coding	3	1492	cagggtcagcgagtaggc	11	25
27914	Coding	3	1589	gcacgaagcagaggtggg	53	26
27915	Coding	3	1746	ttgacacactgggtgggc	39	27
27916	Coding	3	1851	gggtggcacggcaaacag	19	28
27917	Coding	3	1977	tcaggtttcacaccgctg	34	29
27918	Coding	3	2130	accgcagagatgatggac	2	30
27919	Coding	3	2155	gaccacgaccagcagaat	24	31
27920	Coding	3	2199	ttctgctgccgtcgcttg	25	32
27921	Coding	3	2281	gttgggcatcgctccgct	41	33
27922	Coding	3	2367	cccttgtagactgtgcca	44	34
27923	Coding	3	2535	gtggatgtcaggcagatg	14	35
27924	Coding	3	2677	ccgcacatcctccaggtg	36	36
27925	Coding	3	2765	gcagccgagccagcccga	39	37
27926	Coding	3	2866	actctgggtgggtgaaccg	14	38
27927	Coding	3	3185	ccagcagtgcgcggtaga	59	39
27928	Coding	3	3324	ccgccactcctggtagat	0	40
27929	Coding	3	3408	tcggagccagccccttcg	47	41
27930	Coding	3	3534	ccatcagtctcagagggc	70	42
27931	Coding	3	3664	cctttccagagtggcacc	25	43
27932	Coding	3	3809	cgaaggctgggctgaagg	27	44
27933	Stop Codon	3	3930	ctggttcacactggcacg	60	45
27934	3'UTR	3	3999	tgatgccagcagaagtca	6	46
27935	3'UTR	3	4088	gggaactcaagcaggaag	32	47
27936	3'UTR	3	4148	ccacaaagactcccagt	56	48

As shown in Table 1, SEQ ID NOs 10, 12, 15, 18, 21, 22, 23, 24, 26, 27, 29, 31, 32, 33, 34, 36, 37, 39, 41, 42, 43, 44, 45, 47 and 48 demonstrated at least 20% inhibition of human Her-2 expression in this assay and are therefore preferred. The target sites to which these preferred sequences are complementary are herein referred to as "active sites" and are therefore preferred sites for targeting by compounds of the present invention.

-88-

Example 16

Antisense inhibition of Her-2 expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap.

5 In accordance with the present invention, a second series of oligonucleotides were designed to target different regions of the Her-2 RNA, using published sequences (GenBank accession number X03363, incorporated herein as SEQ ID NO: 3). The oligonucleotides are shown in
 10 Table 2. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 2 are chimeric oligonucleotides ("gapmers") 18 nucleotides in length, composed of a central "gap" region consisting of
 15 ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by four-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine
 20 residues are 5-methylcytidines. The compounds were analyzed for their effect on Her-2 mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments. If present, "N.D." indicates "no data".

25

Table 2

Inhibition of Her-2 mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

30

ISIS #	REGION	TARGET (SEQ ID NO)	TARGET SITE	SEQUENCE	%INHIB	SEQ ID NO
27938	Coding	3	280	ggctctcgggactggcagg	58	10
27939	Coding	3	310	ctggtagaggtggcggag	0	11

-89-

27940	Coding	3	326	gcaccacctggcagccct	10	12
27941	Coding	3	365	tggcattggtgggcaggt	25	13
27942	Coding	3	379	caggaaggacaggctggc	40	14
27943	Coding	3	437	cctgcctcacttggttgt	32	15
27944	Coding	3	501	gccagggcatagttgtcc	71	16
27945	Coding	3	539	gggtggtattgttcagcg	55	17
27946	Coding	3	692	gctgggtgttcttgtgga	67	18
27947	Coding	3	769	gcagcgggagcccttaca	55	19
27948	Coding	3	937	gtggttgaagtggaggca	56	20
27949	Coding	3	1159	gcacttctcacaccgctg	73	21
27950	Coding	3	1190	gaccatagcacactcggg	68	22
27951	Coding	3	1232	cactggtaactgccctca	60	23
27952	Coding	3	1327	ggcagtgttgaggctgg	69	24
27953	Coding	3	1492	cagggtcagcgagtaggc	53	25
27954	Coding	3	1589	gcacgaagcagaggtggg	76	26
27955	Coding	3	1746	ttgacacactgggtgggc	59	27
27956	Coding	3	1851	gggtggcacggcaaacag	37	28
27957	Coding	3	1977	tcaggtttcacaccgctg	65	29
27958	Coding	3	2130	accgcagagatgatggac	63	30
27959	Coding	3	2155	gaccacgaccagcagaat	18	31
27960	Coding	3	2199	ttctgctgccgtcgcttg	43	32
27961	Coding	3	2281	gttgggcatcgctccgct	57	33
27962	Coding	3	2367	cccttgtagactgtgcca	75	34
27963	Coding	3	2535	gtggatgtcaggcagatg	55	35
27964	Coding	3	2677	ccgcacatcctccaggta	62	36
27965	Coding	3	2765	gcagccgagccagcccgga	71	37
27966	Coding	3	2866	actctggtgggtgaaccg	50	38
27967	Coding	3	3185	ccagcagtgcagcggtaga	73	39
27968	Coding	3	3324	ccgccactcctggtagat	37	40
27969	Coding	3	3408	tcggagccagcccttcg	72	41
27970	Coding	3	3534	ccatcagtctcagagggc	84	42
27971	Coding	3	3664	cctttccagagtggcacc	65	43
27972	Coding	3	3809	cgaaggctgggctgaagg	48	44
27973	Stop Codon	3	3930	ctggttcacactggcacg	67	45
27974	3'UTR	3	3999	tgatgccagcagaagtca	62	46
27975	3'UTR	3	4088	gggaactcaagcaggaag	0	47
27976	3'UTR	3	4148	ccacaaagactccccagt	68	48

As shown in Table 2, SEQ ID NOs 21, 30, 29, 28, 27, 26, 25, 24, 32, 22, 14, 20, 19, 18, 17, 16, 15, 13, 10, 23, 33, 48, 46, 45, 44, 43, 42, 40, 39, 38, 37, 34, 36, 35 and 5 41 demonstrated at least 20% inhibition of Her-2 expression in this experiment and are therefore preferred. The target sites to which these preferred sequences are complementary are herein referred to as "active sites" and are therefore

-90-

preferred sites for targeting by compounds of the present invention.

Example 17

5 Western blot analysis of Her-2 protein levels

Western blot analysis (immunoblot analysis) is carried out using standard methods. Cells are harvested 16-20 h after oligonucleotide treatment, washed once with PBS, suspended in Laemmli buffer (100 ul/well), boiled for 5
10 minutes and loaded on a 16% SDS-PAGE gel. Gels are run for 1.5 hours at 150 V, and transferred to membrane for western blotting. Appropriate primary antibody directed to Her-2 is used, with a radiolabelled or fluorescently labeled secondary antibody directed against the primary antibody
15 species. Bands are visualized using a PHOSPHORIMAGER™ (Molecular Dynamics, Sunnyvale CA).

-91-

What is claimed is:

1. A compound 8 to 50 nucleobases in length
5 targeted to a nucleic acid molecule encoding Her-2, wherein
said compound specifically hybridizes with and inhibits the
expression of Her-2 and has a sequence comprising SEQ ID
NO: 10, 12, 15, 18, 21, 22, 23, 24, 26, 27, 29, 31, 32, 33,
34, 36, 37, 39, 41, 42, 43, 44, 45, 47, 48, 30, 28, 25, 14,
10 20, 19, 17, 16, 13, 46, 40, 38 or 35.
2. The compound of claim 1 which is an antisense
oligonucleotide.
3. The compound of claim 2 wherein the antisense
oligonucleotide has a sequence comprising SEQ ID NO: 10,
15 15, 18, 21, 22, 23, 24, 26, 27, 29, 32, 33, 34, 36, 37, 39,
41, 42, 43, 44, 45 or 48.
4. The compound of claim 2 wherein the antisense
oligonucleotide comprises at least one modified
internucleoside linkage.
- 20 5. The compound of claim 4 wherein the modified
internucleoside linkage is a phosphorothioate linkage.
6. The compound of claim 2 wherein the antisense
oligonucleotide comprises at least one modified sugar
moiety.
- 25 7. The compound of claim 6 wherein the modified
sugar moiety is a 2'-O-methoxyethyl sugar moiety.
8. The compound of claim 2 wherein the antisense
oligonucleotide comprises at least one modified nucleobase.
9. The compound of claim 8 wherein the modified
30 nucleobase is a 5-methylcytosine.
10. The compound of claim 2 wherein the antisense
oligonucleotide is a chimeric oligonucleotide.
11. The compound of claim 1 at least 8 nucleobases
in length which specifically hybridizes with at least an 8-

-92-

nucleobase portion of an active site on a nucleic acid molecule encoding Her-2.

12. A composition comprising the compound of claim 1 and a pharmaceutically acceptable carrier or diluent.

5 13. The composition of claim 12 further comprising a colloidal dispersion system.

14. The composition of claim 12 wherein the compound is an antisense oligonucleotide.

10 15. A method of inhibiting the expression of Her-2 in cells or tissues comprising contacting said cells or tissues with the compound of claim 1 so that expression of Her-2 is inhibited.

15 16. A method of treating a human having a disease or condition associated with Her-2 comprising administering to said animal a therapeutically or prophylactically effective amount of the compound of claim 1 so that expression of Her-2 is inhibited.

17. The method of claim 16 wherein the disease or condition is a hyperproliferative condition.

20 18. The method of claim 17 wherein the hyperproliferative condition is cancer.

19. The method of claim 18 wherein the cancer is lung, breast, gastric, esophageal, colon, bladder, salivary, neural, or cardiac.

-1-

SEQUENCE LISTING

<110> Isis Pharmaceuticals, Inc.
C. Frank Bennett
Lex M. Cowser

<120> ANTISENSE MODULATION OF HER-2 EXPRESSION

<130> RTSP-0183

<150> 09/663,634

<151> 2000-09-15

<160> 48

<210> 1

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 1

tccgtcatcg ctctcaggg

20

<210> 2

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 2

atgcattctg cccccaagga

20

<210> 3

<211> 4473

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (175)...(3942)

<400> 3

aagggagggt aacctggcc cctttggtcg gggccccggg cagccgcgcg ccccttccca 60

cggggccctt tactgcgcgc cgcgcccggc cccaccct cgcagcacc cgcgccccgc 120

-2-

gccctcccag ccgggtccag ccggagccat ggggccggag ccgcagtgag cacc atg	177
Met	
1	
gag ctg gcg gcc ttg tgc cgc tgg ggg ctc ctc ctc gcc ctc ttg ccc	225
Glu Leu Ala Ala Leu Cys Arg Trp Gly Leu Leu Leu Ala Leu Leu Pro	
5 10 15	
ccc gga gcc gcg agc acc caa gtg tgc acc ggc aca gac atg aag ctg	273
Pro Gly Ala Ala Ser Thr Gln Val Cys Thr Gly Thr Asp Met Lys Leu	
20 25 30	
cgg ctc cct gcc agt ccc gag acc cac ctg gac atg ctc cgc cac ctc	321
Arg Leu Pro Ala Ser Pro Gly Thr His Leu Asp Met Leu Arg His Leu	
35 40 45	
tac cag ggc tgc cag gtg gtg cag gga aac ctg gaa ctc acc tac ctg	369
Tyr Gln Gly Cys Gln Val Val Gln Gly Asn Leu Glu Leu Thr Tyr Leu	
50 55 60 65	
ccc acc aat gcc agc ctg tcc ttc ctg cag gat atc cag gag gtg cag	417
Pro Thr Asn Ala Ser Leu Ser Phe Leu Gln Asp Ile Gln Glu Val Gln	
70 75 80	
ggc tac gtg ctc atc gct cac aac caa gtg agg cag gtc cca ctg cag	465
Gly Tyr Val Leu Ile Ala His Asn Gln Val Arg Gln Val Pro Leu Gln	
85 90 95	
agg ctg cgg att gtg cga ggc acc cag ctc ttt gag gac aac tat gcc	513
Arg Leu Arg Ile Val Arg Gly Thr Gln Leu Phe Glu Asp Asn Tyr Ala	
100 105 110	
ctg gcc gtg cta gac aat gga gac ccg ctg aac aat acc acc cct gtc	561
Leu Ala Val Leu Asp Asn Gly Asp Pro Leu Asn Asn Thr Thr Pro Val	
115 120 125	
aca ggg gcc tcc cca gga ggc ctg cgg gag ctg cag ctt cga agc ctc	609
Thr Gly Ala Ser Pro Gly Gly Leu Arg Glu Leu Gln Leu Arg Ser Leu	
130 135 140 145	
aca gag atc ttg aaa gga ggg gtc ttg atc cag cgg aac ccc cag ctc	657
Thr Glu Ile Leu Lys Gly Gly Val Leu Ile Gln Arg Asn Pro Gln Leu	
150 155 160	
tgc tac cag gac acg att ttg tgg aag gac atc ttc cac aag aac aac	705
Cys Tyr Gln Asp Thr Ile Leu Trp Lys Asp Ile Phe His Lys Asn Asn	
165 170 175	
cag ctg gct ctc aca ctg ata gac acc aac cgc tct cgg gcc tgc cac	753
Gln Leu Ala Leu Thr Leu Ile Asp Thr Asn Arg Ser Arg Ala Cys His	
180 185 190	

-3-

ccc tgt tct ccg atg tgt aag ggc tcc cgc tgc tgg gga gag agt tct Pro Cys Ser Pro Met Cys Lys Gly Ser Arg Cys Trp Gly Glu Ser Ser 195 200 205	801
gag gat tgt cag agc ctg acg cgc act gtc tgt gcc ggt ggc tgt gcc Glu Asp Cys Gln Ser Leu Thr Arg Thr Val Cys Ala Gly Gly Cys Ala 210 215 220 225	849
cgc tgc aag ggg cca ctg ccc act gac tgc tgc cat gag cag tgt gct Arg Cys Lys Gly Pro Leu Pro Thr Asp Cys Cys His Glu Gln Cys Ala 230 235 240	897
gcc ggc tgc acg ggc ccc aag cac tct gac tgc ctg gcc tgc ctc cac Ala Gly Cys Thr Gly Pro Lys His Ser Asp Cys Leu Ala Cys Leu His 245 250 255	945
ttc aac cac agt ggc atc tgt gag ctg cac tgc cca gcc ctg gtc acc Phe Asn His Ser Gly Ile Cys Glu Leu His Cys Pro Ala Leu Val Thr 260 265 270	993
tac aac aca gac acg ttt gag tcc atg ccc aat ccc gag ggc cgg tat Tyr Asn Thr Asp Thr Phe Glu Ser Met Pro Asn Pro Glu Gly Arg Tyr 275 280 285	1041
aca ttc ggc gcc agc tgt gtg act gcc tgt ccc tac aac tac ctt tct Thr Phe Gly Ala Ser Cys Val Thr Ala Cys Pro Tyr Asn Tyr Leu Ser 290 295 300 305	1089
acg gac gtg gga tcc tgc acc ctc gtc tgc ccc ctg cac aac caa gag Thr Asp Val Gly Ser Cys Thr Leu Val Cys Pro Leu His Asn Gln Glu 310 315 320	1137
gtg aca gca gag gat gga aca cag cgg tgt gag aag tgc agc aag ccc Val Thr Ala Glu Asp Gly Thr Gln Arg Cys Glu Lys Cys Ser Lys Pro 325 330 335	1185
tgt gcc cga gtg tgc tat ggt ctg ggc atg gag cac ttg cga gag gtg Cys Ala Arg Val Cys Tyr Gly Leu Gly Met Glu His Leu Arg Glu Val 340 345 350	1233
agg gca gtt acc agt gcc aat atc cag gag ttt gct ggc tgc aag aag Arg Ala Val Thr Ser Ala Asn Ile Gln Glu Phe Ala Gly Cys Lys Lys 355 360 365	1281
atc ttt ggg agc ctg gca ttt ctg ccg gag agc ttt gat ggg gac cca Ile Phe Gly Ser Leu Ala Phe Leu Pro Glu Ser Phe Asp Gly Asp Pro 370 375 380 385	1329
gcc tcc aac act gcc ccg ctc cag cca gag cag ctc caa gtg ttt gag Ala Ser Asn Thr Ala Pro Leu Gln Pro Glu Gln Leu Gln Val Phe Glu 390 395 400	1377

- 4 -

act ctg gaa gag atc aca ggt tac cta tac atc tca gca tgg ccg gac	1425
Thr Leu Glu Glu Ile Thr Gly Tyr Leu Tyr Ile Ser Ala Trp Pro Asp	
405 410 415	
agc ctg cct gac ctc agc gtc ttc cag aac ctg caa gta atc cgg gga	1473
Ser Leu Pro Asp Leu Ser Val Phe Gln Asn Leu Gln Val Ile Arg Gly	
420 425 430	
cga att ctg cac aat ggc gcc tac tgc ctg acc ctg caa ggg ctg ggc	1521
Arg Ile Leu His Asn Gly Ala Tyr Ser Leu Thr Leu Gln Gly Leu Gly	
435 440 445	
atc agc tgg ctg ggg ctg cgc tca ctg agg gaa ctg ggc agt gga ctg	1569
Ile Ser Trp Leu Gly Leu Arg Ser Leu Arg Glu Leu Gly Ser Gly Leu	
450 455 460 465	
gcc ctc atc cac cat aac acc cac ctc tgc ttc gtg cac acg gtg ccc	1617
Ala Leu Ile His His Asn Thr His Leu Cys Phe Val His Thr Val Pro	
470 475 480	
tgg gac cag ctc ttt cgg aac ccg cac caa gct ctg ctc cac act gcc	1665
Trp Asp Gln Leu Phe Arg Asn Pro His Gln Ala Leu Leu His Thr Ala	
485 490 495	
aac cgg cca gag gac gag tgt gtg ggc gag ggc ctg gcc tgc cac cag	1713
Asn Arg Pro Glu Asp Glu Cys Val Gly Glu Gly Leu Ala Cys His Gln	
500 505 510	
ctg tgc gcc cga ggg cac tgc tgg ggt cca ggg ccc acc cag tgt gtc	1761
Leu Cys Ala Arg Gly His Cys Trp Gly Pro Gly Pro Thr Gln Cys Val	
515 520 525	
aac tgc agc cag ttc ctt cgg ggc cag gag tgc gtg gag gaa tgc cga	1809
Asn Cys Ser Gln Phe Leu Arg Gly Gln Glu Cys Val Glu Glu Cys Arg	
530 535 540 545	
gta ctg cag ggg ctc ccc agg gag tat gtg aat gcc agg cac tgt ttg	1857
Val Leu Gln Gly Leu Pro Arg Glu Tyr Val Asn Ala Arg His Cys Leu	
550 555 560	
ccg tgc cac cct gag tgt cag ccc cag aat ggc tca gtg acc tgt ttt	1905
Pro Cys His Pro Glu Cys Gln Pro Gln Asn Gly Ser Val Thr Cys Phe	
565 570 575	
gga ccg gag gct gac cag tgt gtg gcc tgt gcc cac tat aag gac cct	1953
Gly Pro Glu Ala Asp Gln Cys Val Ala Cys Ala His Tyr Lys Asp Pro	
580 585 590	
ccc ttc tgc gtg gcc cgc tgc ccc agc ggt gtg aaa cct gac ctc tcc	2001
Pro Phe Cys Val Ala Arg Cys Pro Ser Gly Val Lys Pro Asp Leu Ser	
595 600 605	

-5-

tac atg ccc atc tgg aag ttt cca gat gag gag ggc gca tgc cag cct	2049
Tyr Met Pro Ile Trp Lys Phe Pro Asp Glu Glu Gly Ala Cys Gln Pro	
610 615 620 625	
tgc ccc atc aac tgc acc cac tcc tgt gtg gac ctg gat gac aag ggc	2097
Cys Pro Ile Asn Cys Thr His Ser Cys Val Asp Leu Asp Asp Lys Gly	
630 635 640	
tgc ccc gcc gag cag aga gcc agc cct ctg acg tcc atc atc tct gcg	2145
Cys Pro Ala Glu Gln Arg Ala Ser Pro Leu Thr Ser Ile Ile Ser Ala	
645 650 655	
gtg gtt ggc att ctg ctg gtc gtg gtc ttg ggg gtg gtc ttt ggg atc	2193
Val Val Gly Ile Leu Leu Val Val Val Leu Gly Val Val Phe Gly Ile	
660 665 670	
ctc atc aag cga cgg cag cag aag atc cgg aag tac acg atg cgg aga	2241
Leu Ile Lys Arg Arg Gln Gln Lys Ile Arg Lys Tyr Thr Met Arg Arg	
675 680 685	
ctg ctg cag gaa acg gag ctg gtg gag ccg ctg aca cct agc gga gcg	2289
Leu Leu Gln Glu Thr Glu Leu Val Glu Pro Leu Thr Pro Ser Gly Ala	
690 695 700 705	
atg ccc aac cag gcg cag atg cgg atc ctg aaa gag acg gag ctg agg	2337
Met Pro Asn Gln Ala Gln Met Arg Ile Leu Lys Glu Thr Glu Leu Arg	
710 715 720	
aag gtg aag gtg ctt gga tct ggc gct ttt ggc aca gtc tac aag ggc	2385
Lys Val Lys Val Leu Gly Ser Gly Ala Phe Gly Thr Val Tyr Lys Gly	
725 730 735	
atc tgg atc cct gat ggg gag aat gtg aaa att cca gtg gcc atc aaa	2433
Ile Trp Ile Pro Asp Gly Glu Asn Val Lys Ile Pro Val Ala Ile Lys	
740 745 750	
gtg ttg agg gaa aac aca tcc ccc aaa gcc aac aaa gaa atc tta gac	2481
Val Leu Arg Glu Asn Thr Ser Pro Lys Ala Asn Lys Glu Ile Leu Asp	
755 760 765	
gaa gca tac gtg atg gct ggt gtg ggc tcc cca tat gtc tcc cgc ctt	2529
Glu Ala Tyr Val Met Ala Gly Val Gly Ser Pro Tyr Val Ser Arg Leu	
770 775 780 785	
ctg ggc atc tgc ctg aca tcc acg gtg cag ctg gtg aca cag ctt atg	2577
Leu Gly Ile Cys Leu Thr Ser Thr Val Gln Leu Val Thr Gln Leu Met	
790 795 800	
ccc tat ggc tgc ctc tta gac cat gtc cgg gaa aac cgc gga cgc ctg	2625
Pro Tyr Gly Cys Leu Leu Asp His Val Arg Glu Asn Arg Gly Arg Leu	
805 810 815	

- 6 -

ggc tcc cag gac ctg ctg aac tgg tgt atg cag att gcc aag ggg atg Gly Ser Gln Asp Leu Leu Asn Trp Cys Met Gln Ile Ala Lys Gly Met 820 825 830	2673
agc tac ctg gag gat gtg cgg ctc gta cac agg gac ttg gcc gct cgg Ser Tyr Leu Glu Asp Val Arg Leu Val His Arg Asp Leu Ala Ala Arg 835 840 845	2721
aac gtg ctg gtc aag agt ccc aac cat gtc aaa att aca gac ttc ggg Asn Val Leu Val Lys Ser Pro Asn His Val Lys Ile Thr Asp Phe Gly 850 855 860 865	2769
ctg gct cgg ctg ctg gac att gac gag aca gag tac cat gca gat ggg Leu Ala Arg Leu Leu Asp Ile Asp Glu Thr Glu Tyr His Ala Asp Gly 870 875 880	2817
ggc aag gtg ccc atc aag tgg atg gcg ctg gag tcc att ctc cgc cgg Gly Lys Val Pro Ile Lys Trp Met Ala Leu Glu Ser Ile Leu Arg Arg 885 890 895	2865
cgg ttc acc cac cag agt gat gtg tgg agt tat ggt gtg act gtg tgg Arg Phe Thr His Gln Ser Asp Val Trp Ser Tyr Gly Val Thr Val Trp 900 905 910	2913
gag ctg atg act ttt ggg gcc aaa cct tac gat ggg atc cca gcc cgg Glu Leu Met Thr Phe Gly Ala Lys Pro Tyr Asp Gly Ile Pro Ala Arg 915 920 925	2961
gag atc cct gac ctg ctg gaa aag ggg gag cgg ctg ccc cag ccc ccc Glu Ile Pro Asp Leu Leu Glu Lys Gly Glu Arg Leu Pro Gln Pro Pro 930 935 940 945	3009
atc tgc acc att gat gtc tac atg atc atg gtc aaa tgt tgg atg att Ile Cys Thr Ile Asp Val Tyr Met Ile Met Val Lys Cys Trp Met Ile 950 955 960	3057
gac tct gaa tgt cgg cca aga ttc cgg gag ttg gtg tct gaa ttc tcc Asp Ser Glu Cys Arg Pro Arg Phe Arg Glu Leu Val Ser Glu Phe Ser 965 970 975	3105
cgc atg gcc agg gac ccc cag cgc ttt gtg gtc atc cag aat gag gac Arg Met Ala Arg Asp Pro Gln Arg Phe Val Val Ile Gln Asn Glu Asp 980 985 990	3153
ttg ggc cca gcc agt ccc ttg gac agc acc ttc tac cgc tca ctg ctg Leu Gly Pro Ala Ser Pro Leu Asp Ser Thr Phe Tyr Arg Ser Leu Leu 995 1000 1005	3201
gag gac gat gac atg ggg gac ctg gtg gat gct gag gag tat ctg gta Glu Asp Asp Asp Met Gly Asp Leu Val Asp Ala Glu Glu Tyr Leu Val 1010 1015 1020 1025	3249

- 7 -

ccc cag cag ggc ttc ttc tgt cca gac cct gcc ccg ggc gct ggg ggc	3297
Pro Gln Gln Gly Phe Phe Cys Pro Asp Pro Ala Pro Gly Ala Gly Gly	
1030 1035 1040	
atg gtc cac cac agg cac cgc agc tca tct acc agg agt ggc ggt ggg	3345
Met Val His His Arg His Arg Ser Ser Ser Thr Arg Ser Gly Gly Gly	
1045 1050 1055	
gac ctg aca cta ggg ctg gag ccc tct gaa gag gag gcc ccc agg tct	3393
Asp Leu Thr Leu Gly Leu Glu Pro Ser Glu Glu Glu Ala Pro Arg Ser	
1060 1065 1070	
cca ctg gca ccc tcc gaa ggg gct ggc tcc gat gta ttt gat ggt gac	3441
Pro Leu Ala Pro Ser Glu Gly Ala Gly Ser Asp Val Phe Asp Gly Asp	
1075 1080 1085	
ctg gga atg ggg gca gcc aag ggg ctg caa agc ctc ccc aca cat gac	3489
Leu Gly Met Gly Ala Ala Lys Gly Leu Gln Ser Leu Pro Thr His Asp	
1090 1095 1100 1105	
ccc agc cct cta cag cgg tac agt gag gac ccc aca gta ccc ctg ccc	3537
Pro Ser Pro Leu Gln Arg Tyr Ser Glu Asp Pro Thr Val Pro Leu Pro	
1110 1115 1120	
tct gag act gat ggc tac gtt gcc ccc ctg acc tgc agc ccc cag cct	3585
Ser Glu Thr Asp Gly Tyr Val Ala Pro Leu Thr Cys Ser Pro Gln Pro	
1125 1130 1135	
gaa tat gtg aac cag cca gat gtt cgg ccc cag ccc cct tcg ccc cga	3633
Glu Tyr Val Asn Gln Pro Asp Val Arg Pro Gln Pro Pro Ser Pro Arg	
1140 1145 1150	
gag ggc cct ctg cct gct gcc cga cct gct ggt gcc act ctg gaa agg	3681
Glu Gly Pro Leu Pro Ala Ala Arg Pro Ala Gly Ala Thr Leu Glu Arg	
1155 1160 1165	
ccc aag act ctc tcc cca ggg aag aat ggg gtc gtc aaa gac gtt ttt	3729
Pro Lys Thr Leu Ser Pro Gly Lys Asn Gly Val Val Lys Asp Val Phe	
1170 1175 1180 1185	
gcc ttt ggg ggt gcc gtg gag aac ccc gag tac ttg aca ccc cag gga	3777
Ala Phe Gly Gly Ala Val Glu Asn Pro Glu Tyr Leu Thr Pro Gln Gly	
1190 1195 1200	
gga gct gcc cct cag ccc cac cct cct cct gcc ttc agc cca gcc ttc	3825
Gly Ala Ala Pro Gln Pro His Pro Pro Pro Ala Phe Ser Pro Ala Phe	
1205 1210 1215	
gac aac ctc tat tac tgg gac cag gac cca cca gag cgg ggg gct cca	3873
Asp Asn Leu Tyr Tyr Trp Asp Gln Asp Pro Pro Glu Arg Gly Ala Pro	
1220 1225 1230	

- 8 -

ccc agc acc ttc aaa ggg aca cct acg gca gag aac cca gag tac ctg 3921
 Pro Ser Thr Phe Lys Gly Thr Pro Thr Ala Glu Asn Pro Glu Tyr Leu
 1235 1240 1245

ggt ctg gac gtg cca gtg tga accagaaggc caagtccgca gaagccctga 3972
 Gly Leu Asp Val Pro Val
 1250 1255

tgtgtcctca gggagcaggg aaggcctgac ttctgctggc atcaagaggt gggagggccc 4032

tccgaccact tccaggggaa cctgccatgc caggaaacctg tcctaaggaa ccttccttcc 4092

tgcttgagtt cccagatggc tggaaggggt ccagcctcgt tggaagagga acagcactgg 4152

ggagtctttg tggattctga ggccctgccc aatgagactc taggggtccag tggatgccac 4212

agcccagctt ggccctttcc ttccagatcc tgggtactga aagccttagg gaagctggcc 4272

tgagagggga agcggcccta agggagtgtc taagaacaaa agcgacccat tcagagactg 4332

tccctgaaac ctagtactgc ccccatgag gaaggaaacag caatgggtgtc agtatccagg 4392

ctttgtacag agtgcctttc tgtttagttt ttactttttt tgttttgttt ttttaaagat 4452

gaaataaaga cccaggggga g 4473

<210> 4

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 4

gtgacagcag aggatggaac ac

22

<210> 5

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 5

atgcccagac catagcacac t

21

- 9 -

<210> 6
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Probe

<400> 6
cgggtgtgaga agtgcagcaa gccc

24

<210> 7
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 7
caacggattt ggtcgtattg g

21

<210> 8
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 8
ggcaacaata tccactttac cagagt

26

<210> 9
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Probe

<400> 9
cgcttggtca ccagggtgc t

21

<210> 10
<211> 18
<212> DNA
<213> Artificial Sequence

-10-

<220>
<223> Antisense Oligonucleotide

<400> 10
ggtctcggga ctggcagg

18

<210> 10
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 10
ggtctcggga ctggcagg

18

<210> 11
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 11
ctggtagagg tggcggag

18

<210> 12
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 12
gcaccacctg gcagccct

18

<210> 13
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 13
tggcattggt gggcaggt

18

-11-

<210> 14
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 14
caggaaggac aggctggc 18

<210> 15
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 15
cctgcctcac ttggttgt 18

<210> 16
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 16
gccagggcat agttgtcc 18

<210> 17
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 17
gggtggtatt gttagcg 18

<210> 18
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

-12-

<400> 18
gctggttggt cttgtgga 18

<210> 19
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 19
gcagcgggag cccttaca 18

<210> 20
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 20
gtggttgaag tggaggca 18

<210> 21
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 21
gcacttctca caccgctg 18

<210> 21
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 21
gcacttctca caccgctg 18

<210> 22
<211> 18
<212> DNA
<213> Artificial Sequence

-13-

<223> Antisense Oligonucleotide

<400> 22

gaccatagca cactcggg

18

<210> 22

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 22

gaccatagca cactcggg

18

<210> 23

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 23

cactggtaac tgccctca

18

<210> 23

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 23

cactggtaac tgccctca

18

<210> 24

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 24

ggcagtgttg gaggctgg

18

<210> 24

-14-

<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 24
ggcagtgttg gaggtgg 18

<210> 25
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 25
cagggtcagc gagtaggc 18

<210> 25
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 25
cagggtcagc gagtaggc 18

<210> 26
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 26
gcacgaagca gaggtggg 18

<210> 26
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 26
gcacgaagca gaggtggg 18

-15-

<210> 27
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 27
ttgacacact gggtgggc 18

<210> 27
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 27
ttgacacact gggtgggc 18

<210> 28
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 28
gggtggcacg gcaaacag 18

<210> 28
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 28
gggtggcacg gcaaacag 18

<210> 29
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

-16-

<400> 29
tcaggtttca caccgctg 18

<210> 30
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 30
accgcagaga tgatggac 18

<210> 31
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 31
gaccacgacc agcagaat 18

<210> 32
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 32
ttctgctgcc gtcgcttg 18

<210> 33
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 33
gttgggcatc gctccgct 18

<210> 34
<211> 18
<212> DNA

-17-

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 34

cccttgtaga ctgtgccca

18

<210> 35

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 35

gtggatgtca ggcagatg

18

<210> 36

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 36

ccgcacatcc tccaggta

18

<210> 37

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 37

gcagccgagc cagcccga

18

<210> 38

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 38

actctggtgg gtgaaccg

18

-18-

<210> 39
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 39
ccagcagtga gcggtaga 18

<210> 40
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 40
ccgccactcc tggtagat 18

<210> 41
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 41
tcggagccag ccccttcg 18

<210> 42
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 42
ccatcagtct cagagggc 18

<210> 43
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 43

-19-

cctttccaga gtggcacc 18

<210> 44
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 44
cgaaggctgg gctgaagg 18

<210> 45
<211> 18
<212> DNA
<213> Artificial Sequence ,

<223> Antisense Oligonucleotide

<400> 45
ctggttcaca ctggcacg 18

<210> 46
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 46
tgatgccagc agaagtca 18

<210> 47
<211> 18
<212> DNA
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 47
gggaactcaa gcaggaag 18

<210> 48
<211> 18
<212> DNA
<213> Artificial Sequence

-20-

<223> Antisense Oligonucleotide

<400> 48

ccacaaagac tccccagt

18

16

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/US01/28572

A. CLASSIFICATION OF SUBJECT MATTER IPC(7) : Please See Extra Sheet. US CL : Please See Extra Sheet. According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 436/6, 91.1, 91.3, 325, 375; 536/23.1, 23.2, 24.5, 24.3, 24.31, 24.33; 514/44 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) biosis, medline, caplus, lifesci, embase, uspatfull		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y	WO 95/17507 A1(BIOGNOSTIK GELLSCHAFT FUR BIOMOLEKULARE DIAGNOSTIK MBH) 29 June 1995 (29.06.95), see especially SEQ ID NO:20 and 34.	1-5, 11-15 -- 6-10
X -- Y	WO 99/57309 A1 (DAKO A/S) 11 November 1999 (11.11.99), see especially SEQ ID NO:67.	1, 2, 4, 11 -- 3, 5-10, 12-15
X -- Y	WO 94/09022 A1 (ONCOR, INC.) 28 April 1994 (28.04.94), see especially page 22, SEQ ID NO:2.	1, 2, 11 -- 3-10, 12-15
X -- Y	US 5,856,089 A (WANG ET AL.) 05 January 1999 (05-01-99), see especially SEQ ID NO:2.	1, 2, 11 -- 3-10, 12-15
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:		
"A"	document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E"	earlier document published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means	"A" document member of the same patent family
"P"	document published prior to the international filing date but later than the priority date claimed	
Date of the actual completion of the international search 15 NOVEMBER 2001		Date of mailing of the international search report 09 JAN 2002
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer KARENA LACOURCIERE Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US01/28572

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,599,704 A (THOMPSON ET AL.) 04 February 1997 (04-2-97), see especially SEQ ID NO: 12, 28, 31, 41, 99 and 2.	1-15
Y	US 5,968,748 A (BENNETT ET AL.) 19 October 1999 (19-10-99), see entire document.	1-15
Y	WO 98/20168 A1 (DUKE UNIVERSITY) 14 May 1998 (14-05-98), see entire document.	1-15
Y	ZAIA ET AL. Status of Ribozyme and Antisense-Based Developmental Approaches for Anti-HIV-1 Therapy. Annals of the New York Academy of Sciences. January 1992, Vol. 660, pages 95-106, see especially page 95.	1-15
X	EP 0 856 579 A1 (BIOGNOSTIK GESELLSCHAFT FUR	1, 2, 11
--	BIOMOLEKULARE DIAGNOSTIK MBH) 05 August 1998 (05-	--
Y	08-98), see entire document.	3-10, 12-15

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US01/28572

A. CLASSIFICATION OF SUBJECT MATTER:
IPC (7):

C07H 21/02, 21/04; A61K 48/00; C12N 15/85, 15/86; C12Q 1/68; C12P 19/34

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

435/6, 91.1, 91.3, 325, 375; 536/23.1, 23.2, 24.5, 24.5, 24.51, 24.53; 514/44